c feeceeie.c

2nd World Conference on

Industrial Chemistry and Water Treatment

May 22-23, 2017 Las Vegas, USA

& ORQLQJ FKDUDFWHUL]DWLRQ DQG VDFFKDUL; FDW*TheFinQtoga*QDO\V petrophila in a mesophilic host

Ikram-ul-Haq and Fatima Akram Government College University, Pakistan

Production of bioethanol has received much attention in recent years and many countries have made large investments in infrastructure, process development and production facilities. Energy crisis are the leading economic constrains in developed as well as in developing countries. With the exhaustion of nonrenewable resources at an exponential rate, the need to develop alternative renewable sources which can be both cost e ective, environmental friendly and high in yield is the need of time. Recently, the increasing demand of energy has strongly stimulated the research on conversion of lignocellulosic plant biomass by the action of cellulases enzymes into reducing sugars, for the subsequent production of bioethanol. Endoglucanases are mainly responsible for hydrolyzing the internal glycosidic bond to decrease the length of the cellulose chains. Obtaining e cient and thermostable endoglucanase has become the goal of much research worldwide. erefore, our research work was focused to search for new resources of endoglucanases, which was thermostable and with high catalytic e ciency. e article focuses on the thermotolerant endo-1,4- -glucanase gene, of ermotoga petrophila RKU-1, was cloned and over-expressed in E. coli strain BL21 codon plus for its potential usage for the hydrolysis of lignocellulosic biomass and in di erent industrial ermostable endoglucanase can be used simultaneously and directly in the sacchari cation procedure without applications. a pre-cooling process of biomass. Puri ed enzyme was optimally active with 530 Umg-1 of speci c activity against CMC at pH 6.0 and 95 °C, which has exhibited a half- life (t_{1/2}) of 6.6 min even at temperature as high as 97 °C and stable up to 8 hours at 80 °C. e recombinant enzyme sacchari ed pre-treated wheat straw and baggase to 3.32% and 3.2%, respectively a er 6 h incubation at 85 °C. Its thermostability, resistance to heavy metal ions and high speci c activity make endoglucanase a potential and promising candidate for various industrial applications such as in textile industry (in bio stoning and bio nishing), in animal feed production, in processing of beer and fruit juice, in biomass hydrolysis (bioethanol production) and in plant oil, detergent, pulp and paper industry.