

JOINT EVENT

5th World Conference on **Climate Change**

&

16th Annual Meeting onOctober 04-06, 2018
London, UK**Environmental Toxicology and Biological Systems****Antarctic marine biodiversity and climate change**

Human culture and food security rely on the ecosystem services provided by historic patterns of biodiversity. We therefore need to understand the factors that determine where species can and cannot live, and the impact of both natural and anthropogenic variation. Such predictions require an understanding of the mechanisms underlying species range limits, and how they are linked to climate. The Southern Ocean offers a “natural laboratory” for testing the evolutionary and physiological capacity of species in response to their environment. Its isolation has resulted in high levels of endemism and the lack of indigenous humans means that the environment is close to pristine. It is a constantly cold ocean but with large seasonal variation in light levels, primary productivity and pH. Animals living in the Southern Ocean have several physiological adaptations for life in the cold, including natural antifreeze, increased mitochondrial densities and the ability to grow to a large size. Life in the extreme cold has also resulted in a reduced ability to cope with warming. The activity limits for limpets and clams are only 1 to 2°C above current maximum summer temperatures. Comparisons of long-term oceanographic and reproductive data-sets have shown that one of the strongest signals affecting inter-annual variability in reproduction is El Niño, which causes dramatic changes in the coastal system. In addition to this understanding, the Western Antarctic Peninsula has been one of the fastest warming regions, resulting in massive changes in the cryosphere. The reduction in the duration of winter sea ice, an increase in energy transfer from the atmosphere and the increase in iceberg scour has resulted in dramatic changes in benthic communities. Findings from the Antarctic have taught us much about the evolution of physiological capacity and the evolution of marine communities across latitudes.

Recent Publications

1. Ashton G V, Morley S A, Barnes D K A, Clark M S and Peck L S (2017) Warming by 1°C drives species and assemblage level responses in Antarctica's marine shallows. *Current Biology* 27(17):2698–2705.
2. Watson S A, Morley S A and Peck L S (2017) Latitudinal trends in shell production