

# Editorial Board

Dr. Ingrid Štěrbová


Faculty of Fisheries & Protection of Waters

Institute of Aquaculture

University of South Bohemia

Czech Republic

Tel. 004-20-777-248-351



Dr. Sabine Sampels, PhD studied Food Chemistry at Rheinische Friedrich  
Wilhlems Universität, Bonn, Germany and received her PhD

**s r n r s s**

Aquaculture and effects of consumption of fish on human health.

Effect of animal nutrition on fish and meat composition.

Fish and fish products assessment .

Lipids and lipid metabolism in animal products and human nutrition.

Shelf life, oxidation and antioxidants in fish and meat.

## Abstracts

papers Zajíc T., Mráz J., 2014, Effect of frying fat and preparation on carp (*Cyprinus carpio*) fillet lipid composition and oxidation. Czech Journal of Food Sciences, ISSN: 1212-1800.

papers Levý, E., Mráz, J., Vejsada, P., Zajíc, T., 2014, Kvalita a gastronomie ryb a rybích výrobků, Jiho česká univerzita v českých Budějovicích, Fakulta rybářství a ochrany vod, Záměstí 728/II, 389 25 Vodňany, p. 252, ISBN 978-80-87437-85-8 (Czech book)

Zajic, T., Mraz, J., papers, Pickova, J., 2013, Fillet quality changes as a result of purging of common carp (*Cyprinus carpio* L.) with special regard to weight loss and lipid profile. Aquaculture, 400-401, 111-119, ISSN: 0044-8486.

Popelář, M., 2013, Oxidation and Antioxidants in Fish and Meat from Farm to Fork. In: Muzzalupo, I. (Ed.), Food industry. InTech, Rijeka, Croatia, pp. 115–144. ISBN 980-953-307-860-6. (Book chapter)

Zajíc, T., Mráz, J., Popelář, M., Pickova, J., 2013. Aplikace technologie finishing feeding do chovu ryb v praktických podmínkách českého rybářství. Edice Metodik, FROV JU, Vodňany, 2013, p. 38 (Czech certified methodology)

Popelář, M. (2013) Processing and Preparation-Two Key Issues to Increase and Preserve Nutritional Value of Fish and Meat Products. J Fisheries Livest Prod 1: 1

Popelář, M., Pickova, J., 2011 Comparison of two different methods for the separation of lipid classes and fatty acid methylation in reindeer and fish muscle, Food Chemistry, 128 811–819.

Pickova, J., Högberg, A., Neil, M., Fatty acid transfer from sow to piglet differs for different polyunsaturated fatty acids (PUFA), *Physiological Research*, 60.

Metabolism and interaction of fatty acids, antioxidants, and biological active substances in animal muscle and consequent quality of animal foods. In: *Lipids: Categories, Biological Functions and Metabolims, Nutrition and Health*; Editor: Columbus F.; Nova Science Publishers, New York, pp. 45-70.

Pickova, P., Berntsen, M., Minor components in Fish Oil and Alternative Oils with Potential Physiological Effect. In: *Fish Oil Replacement and Alternative Lipid Sources in Aquaculture Diets*. Editors: Turchini, G.M.; Ng, W.K.; Tocher, D.R.; CRC Press, Taylor and Francis Group, p. 351–372.

**p |s** Turner, T., Östström Å., Pickova, J., **☛** Metabolism of  $\alpha$ -linolenic- and eicosapentaenoic acid from linseed and algae respectively in reindeer muscle, *Acta Agriculturae Scandinavica A*, 60, 175-186.

**p |s** Åsli, M., Mørkøre, T., **☛** Storage stability and

**p |s** Strandvik B., Pickova J, 2009, Processed Animal Products with Emphasis on Polyunsaturated Fatty Acid Content. European Journal of Lipid Science and Technology., Vol. 111, 481-488.

Strandvik, B., Eriksson, S., Garemo, M., Palsdottir, V., **p |s** .., Pickova, J., **¶ 8** Is the relatively low intake of n-3 fatty acids in Western diet contributing to the obesity epidemics? Lipid Technology, Vol. 20, Nr.3, 2-4.

Svetlana A. Lanina S.A., Toledo P., **p |s** Kamal-Eldin A., Jastrebova J.A., **¶** Comparison of reversed-phase liquid chromatography–mass spectrometry with electrospray and atmospheric pressure chemical ionization for analysis of dietary tocopherols; Journal of Chromatography A, 1157 (2007) 159–170

**p |s** Pickova, J. & Wiklund, E., **¶ 6** Influence of Diet on Fatty Acids and Tocopherols in M. Longissimus Dorsi from Reindeer; Lipids, Vol. 41, no. 5 , 463-472



Wiklund, E., **Þjörðsson**, Manley, T.R., Pickova, J. & Littlejohn, R. P., 2006, Effects of feeding regimen and chilled storage on water holding capacity, colour stability, pigment content and oxidation in red deer (*Cervus elaphus*) meat; *Journal of the Science of Food and Agriculture*, 86, 98-106.

**Þjörðsson** Pickova, J. & Wiklund, E., 2005. Influence of production system, age and sex on carcass parameters and some biochemical meat quality characteristics of reindeer (*Rangifer tarandus tarandus* L.); *Rangifer*, 25 (2): 85-96

**Þjörðsson** 2005. Fatty Acids and antioxidants in reindeer and red deer - emphasis on animal nutrition and consequent meat quality. *Acta Universitatis Agriculturae Suecia*, 2005:31 Swedish University of Agriculture Uppsala, Sweden.

Wiklund, E., Pickova, J., **p |s** ., & Lundström, K. 2001. Fatty acid composition in *M. longissimus lumborum*, ultimate muscle pH values and carcass parameters in reindeer (*Rangifer tarandus tarandus* L) grazed on natural pasture or fed a commercial feed mixture; *Meat Science*, 58, 293-298.

**p |s** Pickova, J. & Wiklund, E., 2004. Fatty acids, antioxidants and oxidation stability of processed reindeer meat; *Meat Science*, 67, 523-532

Wiklund, E., Johansson, L., **p |s** and Malmfors, G. 2004. Tasty and healthy meat from the North. *Food Science Central*. Available at <http://www.foodsciencecentral.com/library.html#ifis/13583>

# AqA A Ar

Aquaculture is the farming of aquatic organisms such as fish, prawns, molluscs, and aquatic plants related directly or indirectly to human consumption.

Over recent years, world seafood consumption has risen and overall production has increased. However, wild-caught production has remained stable, Aquaculture has therefore become increasingly important in meeting local and global demand for seafood.

Stock restoration or "enhancement" is a form of aquaculture in which hatchery fish and shellfish are released into the wild to rebuild wild populations or coastal habitats such as oyster reefs

Aquaculture also includes the production of ornamental fish for the aquarium trade, and growing plant species used in a range of food, pharmaceutical, nutritional, and biotechnology products.

Marine aquaculture refers to the culturing of species that live in the ocean

U.S. marine aquaculture primarily produces oysters, clams, mussels, shrimp, and salmon as well as lesser amounts of cod, moi, yellowtail, barramundi, seabass, and seabream.

Marine aquaculture can take place in the ocean (that is, in cages, on the seafloor, or suspended in the water column) or in on-land, manmade systems such as ponds or tanks.

# Freshwater Aquaculture

Freshwater aquaculture produces species that are native to rivers, lakes, and streams.

U.S. freshwater aquaculture is dominated by catfish but also produces trout, tilapia, and bass.

Freshwater aquaculture takes place primarily in ponds and in on-land, manmade systems such as recirculating aquaculture systems



ro **A** non

**Where are the best places to put an aquaculture farm?**

There are specific criteria that must be considered when investigating a potential site for aquaculture. These include water access, topography, climate, soil type, and proximity to markets, support and infrastructure.





## T AGE T EE Appeal

The applicant has the right to appeal any decision for refusal, or any conditions attached to an approval.

The Decision is usually made under delegated powers by a local planning officer. If a statutory consultee has objected, the application will be determined by the council committee instead of being delegated to a planning officer, this has a bearing on the route of appeal by the applicant



# Controlling diseases in aquaculture systems

**Species-specific** : Many fish diseases are species-specific and are brought on by different conditions. For example, poor quality water affects trout and tilapia in different ways.

Parasitic infections such as *Ichthyophthirius multifiliis* (ich) and *costia* (white spot diseases) are often opportunistic and attack weakened fish.

**Fungal infections and sensitivity** Fungus may also develop on tilapia exposed to low temperatures. It is of little use to treat the fungus, as it is a secondary affliction – the water must be warmed. Some fish strains are more sensitive than others

**Algae control : Ultraviolet (UV) sterilization is a useful tool in RAS disease control, both to eliminate pathogens and for uni-cellular algae control.**


**If a system is carrying a high nitrate load, green water will almost certainly be a problem, resulting in fish and algae competing for oxygen.**

**UV lamps of sufficiently high output and used for extended periods can control algae very effectively. These are fitted in the filtration system behind the pumps, before the water is returned to the fish tanks. These UV lamps result in clear, pathogen-free water**

## Fish breeding do's & don'ts

If you breed two genetically closely-related individuals of the same species, you automatically lose genetic variability but the golden rule is to breed livestock to improve their attributes for aquaculture, and not breed inbred, low quality runts.

The frequent inbreeding of siblings (brother/sister or father/daughter) usually has two results. Certain traits become more fixed (be they good or bad) and the general health of the fish deteriorates.



If all the fish in a particular population (thousands of individuals) are fast-growing, with deep body shape and a late onset of maturity, this may be an indicator of high-quality stock for aquaculture.

But that can only be ascertained by an intensive scientific examination of their life history.

Only once that has been documented can one say that a particular population, **ost ai**, is of high quality, and even then these attributes may not be as well expressed under

## Fish Farming Jobs

The first method is the cage system which use cages that are placed in lakes, ponds and oceans that contain the fish. This method is also widely referred to as off-shore cultivation. Fish are kept in the cage like structures and are **artificially fed** and harvested.

The second method is irrigation ditch or pond systems for raising fish. This basic requirement for this method is to have a ditch or a pond that holds water. This is a unique system because at a small level, fish are artificially fed and the waste produced from the fish is then used to fertilize **farmer's** fields.

The third method of fish farming is called composite fish culture which is a type of fish farming that allows both local fish species and imported fish species to coexist in the same pond. The number of species depends, but it is sometimes upwards of six fish species in a single pond.

The fourth method of fish farming is called integrated recycling systems **which is considered the largest scale method of pure fish** farming. This approach uses large plastic tanks that are placed inside a greenhouse

The last type of fish farming method is called classic fry farming this **method is also known as flow through system**. This is the spot fish species are raised from eggs and are put in streams and released.



**Thank You..!**