

" \$PNQBSBUJWF " TTFTTNFOU PG UIF %JBHOPT  
\$JUSVMMJOBUE 1FQUJEF "OUJCPEJFT BOE 3I  
" SUI SJUJT

Fariba Binesh<sup>1</sup>, Hossein Soleimani Salehabadi<sup>2</sup>, Nasim Behniafard<sup>3\*</sup>, Kian Ranginkaman<sup>4</sup> and Nasrin Behniafard<sup>5</sup>


<sup>1</sup>Department of Pathology, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

<sup>2</sup>Assistant Professor of Internal Medicine, rheumatologist, Department of Internal Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

<sup>3</sup>Researcher of Research and Development Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

<sup>4</sup>General Practitioners, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

<sup>5</sup>Assistant Professor of Pathology, Sub Specialist in Allergy and Ch PractnScienImmun, Sub Sp



\*Corresponding author:

Regarding the anti-CCP titer, 96% of cases with titer higher than 201 (201-1500) had definite RA whereas, 55.4% of patients with titer

RA.

to the symptom duration of less than one year in more than 60% of our patients. Nevertheless, in this study it was interesting to evaluate

