Research Article Open Access

A Low Glycemic Index Diet Combined with an Aerobic-Resistance Exercise Program Reduces Risk Factors Associated with the Metabolic Syndrome

Kendra E Brett [1] and Kelly A Meckling [2*]

[1]Department of Human Kinetics, University of Ottawa, Healthy Active Living and Obesity Research Group, Children's Hospital of Eastern Ontario, Ottawa, ON, Canada
[2]Department of Human Health and Nutritional Sciences, University of Guelph, Guelph, ON, Canada

Keywords: Metabolic syndrome; Low glycemic index; Obesity; Cardiovascular disease

Introduction

*Corresponding author: Kelly A Meckling, Department of Human Health and Nutritional Sciences, University of Guelph, Guelph, ON, Canada, N1G2W1, Tel: 519-824-4120 ext: 53742; Fax: 519-763-5902; E-mail: kmecklin@uoguelph.ca

Received June 27, 2012; Accepted August 04, 2012; Published August 09, 2012

Citation: Brett KE, Meckling KA (2012) A Low Glycemic Index Diet Combined with an Aerobic-Resistance Exercise Program Reduces Risk Factors Associated with the Metabolic Syndrome. J Obes Wt Loss Ther 2:142. doi:10.4172/2165-7904.1000142

Copyright: © 2012 Brett KE, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

advertisements in the newspaper and community. e study was approved by the University of Guelph Research Ethics Board, and all subjects gave informed written consent. Inclusion criteria were as follows: 18-60 years of age, BMI 27 kg/m² and meeting the 2005 IDF de nition of MetS. Individuals were excluded if they were taking any medications known to a ect blood pressure, blood glucose, blood lipids, or weight. While we recognized that a large proportion of the population for whom this dietary strategy may be e cacious, may be on these medications, we wanted to avoid the confounding of medication use in this pilot study. Subjects who previously had stroke, myocardial infarction or other major DVD events were also excluded. e intervention consisted of a 12 week diet and exercise program. Subjects were randomly assigned to one of four diets and all attended three exercise sessions per week. Additional baseline measurements included percent body fat, a 7 day food record and a tness assessment. Daily food records were kept throughout the intervention and periodically 7 day records were analyzed using e Food Processor for Windows

Finnish men: the Kuopio Ischaemic Heart Disease Risk Factor Study. Nutr Metab Cardiovasc Dis 21: 144-149.

35. Ware JH (2003) Interpreting incomplete data in studies of diet and weight loss. N Engl J Med 348: 2136-2137.

36. Wolever TM, Gibbs AL, Mehling C, Chiasson JL, Connelly PW, et al. (2008) The Canadian Trial of Carbohydrates in Diabetes (CCD), a 1-y controlled trial of low-glycemic-index dietary carbohydrate in type 2 diabetes: no effect on glycated hemoglobin but reduction in C-reactive protein. Am J Clin Nutr 87: 114-125.

37. Johnson RK, Goran MI, Poehlman ET (1994) Correlates of over- and underreporting of energy intake in healthy older men and women. Am J Clin Nutr 59: 1286-1290.

38. Lichtman SW, Pisarska K, Berman ER, Pestone M, Dowling H, et al. (1992) Discrepancy between self-reported and actual caloric intake and exercise in obese subjects. N Engl J Med 327: 1893-1898.

39. Sloth B, Krog-Mikkelsen I, Flint A, Tetens I, Björck I, et al. (2004) No difference in body weight decrease between a low-glycemic-index and a high-glycemic-index diet but reduced LDL cholesterol after 10-wk ad libitum intake of the low-glycemic-index diet. Am J Clin Nutr 80: 337-347.

40. Brand-Miller J, Hayne S, Petocz P, Colagiuri S (2003) Low-Glycemic Index Diets in the Management of Diabetes. A meta-analysis of randomized controlled trials. Diabetes Care 26: 2261-2267.

41. Barclay AW, Petocz P, McMillan-Price J, Flood VM, Prvan T, et al. (2008) Glycemic index, glycemic load, and chronic disease risk--a meta-analysis of observational studies. Am J Clin Nutr 87: 627-637.

42. Hare-Bruun H, Nielsen BM, Grau K, Oxlund AL, Heitmann BL (2008) Should glycemic index and glycemic load be considered in dietary recommendations? Nutr Rev 66: 569-590.

43. Janssen I, Fortier A, Hudson R, Ross R (2002) Effects of an energy-restrictive diet with or without exercise on abdominal fat, intermuscular fat, and metabolic risk factors in obese women. Diabetes Care 25: 431-438.

44. Orchard TJ, Temprosa M, Goldberg R, Haffner S, Ratner R, et al. (2005) The effect of metformin and intensive lifestyle intervention on the metabolic syndrome: the Diabetes Prevention Program randomized trial. Ann Intern Med 142: 611-619.