

Adult Cystic Lymphangioma in Posterior Omentum Cavity

Sara Zenjali*, Sanae Jellal, El Aitari Khadija, El Fenni Jamal and Saouab Rachida

Department of Radiology, Mohammed V Military Hospital, Faculty of Medicine and Pharmacy of Rabat, Mohammed V University, Rabat, Morocco

Abstract

A cystic lymphangioma is a non-malignant tumor that arises from the lymphatic vessels, we report the case of a 60-year-old female patient, presenting with chronic abdominal pain, An abdominal computed tomography was performed, revealing a cystic lymphangioma located in the posterior cavity of the omentum.

Keywords: Cystic; Lymphangioma; Imaging; Omentum

Case Report

A cystic lymphangioma is a non-malignant tumor that arises from the lymphatic vessels [1]. The suggested cause is an embryological anomaly, where primary lymphatic cysts do not properly connect with the main lymphatic system [2].

We report the case of a 60-year-old female patient with no significant medical history, presenting with chronic abdominal pain, evolving in the context of afebrile condition, and maintaining general well-being. The clinical examination revealed a mild pain in the epigastric region without a mass syndrome.

An abdominal computed tomography was performed, revealing a hypodense mass of pure liquid density, unilocular, with a thin and regular wall, without septa or vegetations, measuring 51x51x45 mm (APxTxH), located in the posterior cavity of the omentum, respecting adjacent structures (Figure 1).

Over 80% of lymphangiomas are diagnosed in the first year of life, with rare cases in adults. Gender distribution in adulthood is roughly equal [1].

Cystic lymphangiomas (CL) can develop in various anatomical locations, predominantly in the cervical and axial regions. Intra-abdominal cases are rare, comprising less than 5%, and are most commonly located in the mesentery, greater omentum, mesocolon,

and retroperitoneum, with even rarer cases in the posterior cavity of the omentum [1,3].

Cystic lymphangiomas are often asymptomatic, with clinical presentations varying based on size and location. Complications may lead to acute scenarios such as cystic hemorrhage, secondary infections, and obstruction of urinary, biliary tracts, and intestines [3].


Diagnostic imaging involves radiological methods, with ultrasound as the primary screening modality, revealing distinct features. Computed Tomography (CT) scans depict a low-density cyst with a smooth shell, and Magnetic Resonance Imaging (MRI) enhances characterization, showcasing low-signal masses in T2-weighted and high-signal masses in T1-weighted sequences [1,2].

Differential diagnoses for cystic lymphangioma include lymphoma, hydatid cysts, ovarian cysts, digestive duplication, mucinous cystadenomas, and mesenteric cysts [1].

The definitive treatment for abdominal cystic lymphangioma is radical excision, even in asymptomatic cases. However, with increasing tumor size, radical resection becomes more challenging, elevating the risk of local recurrence [3].

References

1. Maghrebi H, Yakoubi C, Beji H, Letaief F, Amin Makni SM, et al. (2022). Intra-abdominal cystic lymphangioma in adults: A case series of 32 patients and literature review. *Ann Med Surg* 81: 104460
2. Jianchun Xiao, Yuming Shao, Shan Zhu, Xiaodong He (2020) Characteristics of adult abdominal cystic Lymphangioma: a single-center Chinese cohort of 12 cases. *Gastroenterol* 20: 244
3. Mohamed Ben Mabrouk, Malek Barka, Waad Farhat, Fathia Harrabi, Mohamed Azzaza, et al. (2015) Intra-Abdominal Cystic Lymphangioma: Report of 21 Cases. *J Cancer Ther* 6: 572.



*Corresponding author: Zenjali Sara, Department of Radiology, Mohammed V Military Hospital, Faculty of Medicine and Pharmacy of Rabat, Mohammed V University, Rabat, Morocco, E-mail: zenjalisara@gmail.com

Received: 02-Jan-2024, Manuscript No: roa-24-125503, **Editor assigned:** 05-Jan-2024, Pre-QC No: roa-24-125503 (PQ), **Reviewed:** 19-Jan-2024, QC No: roa-24-125503, **Revised:** 26-Jan-2024, Manuscript No: roa-24-125503 (R), **Published:** 31-Jan-2024, DOI: 10.4172/2167-7964.1000525

Citation: Zenjali S, Jellal S, Khadija EA, Jamal EF, Rachida S (2024) Adult Cystic Lymphangioma in Posterior Omentum Cavity. *OMICS J Radiol* 13: 525.

Copyright: © 2024 Zenjali S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.