Journal of
Alzheimer's Disease & Parkinsonism
Mini Review Open Access

: Alzheimer's disease; Amyloid-β peptide; Tau protein; Mitochondrial dysfunction; Oxidative stress; Yeast

A **:** Aβ: Amyloid-β Peptide; AD: Alzheimer's Disease; APP: Amyloid Precursor Protein; ATG: Autophagy-related Protein; Drp1p: Dynamin-Related Protein 1; ER: Endoplasmic Reticulum; ERK: Extracellular signal–Regulated Kinase; HIF: Hypoxia-Inducible

sets of neuropathological manifestations, including accumulation of misfolded and aggregated proteins, the extracellular deposition of senile plaques, followed by the intracellular neuro brillary tangles, consisting mainly of aggregates of hyperphosphorylated microtubule associated protein (MAP) referred to as tau [7,8]. MAP tau is a key protein in stabilizing the microtubule architecture that regulates neuron morphology and synaptic strength. In the course of AD hyperphosphorylated tau gets truncated by proteolytic cleavage, being a subject to O-glycosylation, sumoylation, ubiquitinylation, acetylation and some other modi cations [3,4]. When MAP tau is degraded in tauopathic disorders, neuron dysfunction results [9]. Senile plaques are deposits of the amyloid-β peptide (Aβ) produced by the sequential cleavage of β- and γ-secretase at the C terminus of APP. is peptide has an extraordinary ability to undergo conformational changes and is highly amyloidogenic. More than 20 mutations in APP have been linked to familial AD that have altered APP processing with respect to enhanced Aβ generation or aggregation. It is now generally accepted that a progressive accumulation of Aβ aggregates eventually triggers a cascade of cellular changes, including mitochondrial oxidative damage,

***Corresponding author:** Renata Zvyagilskaya, Laboratory of Bioenergetics, Bach Institute of Biochemistry, Research Center of Biotechnology of the Russian Academy of Sciences. 33, bld. 2 Leninsky Ave, Moscow 119071, Russia, Tel: 74959544088; E-mail: renata_z@inbi.ras.ru

Received October 16, 2017; **Accepted** October 23, 2017; **Published** October 30, 2017

Citation: Goleva T, Rogov A, Zvyagilskaya R (2017) Alzheimer's Disease: Molecular Hallmarks and Yeast Models. J Alzheimers Dis Parkinsonism 7: 394. doi: 10.4172/2161-0460.1000394

Copyright: © 2017 Goleva T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

simple, cheap and quick genetic and environmental manipulations, to the large knowledge base and data collections, high-throughput screening technologies and functional genomics that are not possible in humans [93-95], this organism has become a valuable and prevalent eukaryotic model organism to unravel complex and fundamental intracellular mechanisms underlying neurodegeneration [96-104].

Humanized yeasts are also utilized in high-throughput screening of genes that a ect the toxicity of heterologously expressed human proteins, for large-scale chemical screens aiming at the discovery of novel compounds delaying aging or protecting against human age-related diseases.

Several excellent reviews have dealt with speci c aspects of AD, including Aβ toxicity, using this model ([3,4,7,102,105,106] and references therein).

A

However, the fermentation-oriented yeast *S. cerevisiae* (selected for thousands of years for its capacity for alcoholic fermentation) with less abundant mitochondria is hardly a bioenergetic equivalent of high-energy demanding neurons relying almost exclusively on mitochondrial oxidative phosphorylation. *S. cerevisiae* is not the best model organism for studying fragmentation of mitochondria, as they contain small-sized, poorly structured mitochondria. In these respects, *Yarrowia lipolytica*, a non-toxic ascomycetous yeast species having a haploid genome and sexual life cycle and amenable to both classical and molecular genetic techniques [107], an obligate aerobe with the respiratory metabolism closely resembling that of mammalian cells [108,109], vigorously growing on a variety of simple, well de ned and inexpensive media [110], having a long history of use as a producer of heterological proteins [111], possessing an ability to change its morphology (from yeast-like to true mycelium) in response to environmental conditions

15. Cooke MS, Evans MD, Dizdaroglu M, Lunec J (2003) Oxidative DNA damage: Mechanisms, mutation and disease. FASEB J 17: 1195-1214.

16. Lin MT, Beal MF (2006) Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. Nature 443: 787-795.

17. Chmatalova Z, Vyhnalek M, Laczo J, Hort J, Pospisilova R, et al. (2017) Relation of plasma selenium and lipid peroxidation end products in patients with Alzheimer's disease. Physiol Res.

18. Guo L, Tian J, Du H (2017) Mitochondrial dysfunction and synaptic transmission failure in Alzheimer's disease. J Alzheimers Dis 57: 1071-1086.

19. Guo J, Cheng J, North BJ, Wei W (2017) Functional analyses of major cancer-related signaling pathways in Alzheimer's disease etiology. Biochim Biophys Acta 1868: 341-358.

20. Islam MT (2017) Oxidative stress and mitochondrial dysfunction-linked neurodegenerative disorders. Neurol Res 39: 73-82.

21. Kolosova NG, Tyumentsev MA, Muraleva NA, Kiseleva E, Vitovtov AO, et al. (2017) Antioxidant SkQ1 alleviates signs of Alzheimer's disease-like pathology in old OXYS rats by reversing mitochondrial deterioration. Curr Alzheimer Res.

22. Murphy KE, Park JJ (2017) Can co-activation R ' neurvurrptcn

nauutitM E, ɔ , ɔ

62. Delbarba A, Abate G, Prandelli C, Marziano M, Buizza L, et al. (2016) Mitochondrial alterations in peripheral mononuclear blood cells from

Citation: Goleva T, Rogov A, Zvyagilskaya R (2017) Alzheimer's Disease: Molecular Hallmarks and Yeast Models. J Alzheimers Dis Parkinsonism 7: 394. doi: 10.4172/2161-0460.1000394