

An Overview on Renal Pathology of Obesity

Richard Genzen*

Department of Pathology, Memorial Sloan Kettering Cancer Centre, New York, USA

***Corresponding author:** Richard Genzen, Department of Pathology, Memorial Sloan Kettering Cancer Centre, New York, USA, E-mail: Genzen@Richard.vard.edu

Received: 04-Mar-2022, Manuscript No. DPO-22-59987; **Editor assigned:** 07-Mar-2022, PreQC No. DPO-22- 59987 (PQ); **Reviewed:** 18-Mar-2022, QC No. DPO-22- 59987; **Revised:** 22-Mar-2022, Manuscript No. DPO-22- 59987 (R); **Published:** 29-Mar-2022, DOI: 10.4172/2476-2024.1000202

Citation: Genzen R (2022) An Overview on Renal Pathology of Obesity. *Diagnos Pathol Open* 7: 202.

Copyright: © 2022 Genzen R. This is an open-access article distributed under the terms of the

renal mass loss) is very useful in distinguishing these patients from idiopathic FSG, in which a complete nephrotic syndrome is usually present along with nephrotic range proteinuria. The reasons why people with ORG do not develop complete nephrotic syndrome are

unknown, although they may be related to changes in tubular protein management and a very gradual proteinuria growth over time, which is common in ORG and other hyper filtering illnesses.