



... . Pac ... d a bec ... e ... abd ... a a .  
b ed e ... e a d ... a acc ... a e de ... e d, ca  
ec ...

For children, surgery is generally reserved for umbilical hernias  
a:

A e a  
A e ... a e ... a 1/4 3/4 c (1 2 ce ... e e )  
e e

A e a e a d d ... d ... e e e ... e e  
D ... a ... b a e 5

C e a ed b c e e e e

For grown-ups, surgery is generally recommended to avoid  
bec ca , e e ca ... e b ca e a e b e  
bec e a [5].

C ...  
D ... e , a a a ... ade ea e be b ... e  
e a ed ... e e ed ... e abd ... a de e ... , a d e  
e ... e abd ... a a ... ed ... e c ed. I ... ,

A ...  
I ... d ... e ... a ... P ... e ... a d  
e c ... a e e e .

C ...  
e a ... de ca e a ... e a e c c ... e e .

References

- 1. Meier DE, OlaOlorun DA, Omodele RA, Nkor SK, Tarpley JL (2001) "Incidence of indications for repair". World J Surg 25: 645-648.
- 2. Dalenback J, Andersson C, Ribokas D, Rimback G (2013) "Long-term follow-up after elective adult umbilical hernia repair: low recurrence rates also after non-mesh repairs". Hernia 17: 493-497.
- 3. Papagrigoriadis S, Browse DJ, Howard ER (1998) "Incarceration of umbilical