Ayurveda a Boon for Epileptics

Avinash Shankar^{1*}, Amresh Shankar² and Anuradha Shankar²

¹National Institute of Health & Research, Bihar, India ²Centre for Indigenous Medicine & Research, Bihar, India

Abstract

- -----, ..., ..., / · ·
-
- ·····

اردی از داده از به ۲۰۰ در از از به داده از مانه در مرد ویرد به از در از مان از به به معدمین از هار مهم درم مانور دارا از در ورد ما

Obe, a.i.

Grade	Characteristics
I	Complete absence of seizure, improved mental status and physical capacity without any untoward effect and Recurrence.
II	Complete absence of seizure without any untoward effect. Improved physical capabilities but unchanged mental state
III	No response

Table 1: Therapeutic response comparison as per index.



Figure 1: Pie diagram showing sex wise composition of patients.

Age Group	Number of patient			
	Male	Female	Total	%
<5 yr	72	48	120	2.6
5-10 yr	414	330	744	15.97
10-15 yr	480	320	800	17.17
15-20 yr	610	480	1090	23.4
20-25 yr	320	186	506	10.86
25-30 yr	318	236	554	11.89
30-35 yr	276	164	440	9.44
35-40 yr	190	114	304	6.52

Table 2: Distribution of patients as per age and sex.

	30000	
I. Bar Diagram	23500	
a) New Cases - 2580	2000 - 1767	
b) Chronic Cass 1767		
c) Old resistant Case 21311		
	52 2 2 11 J 2 11	
	a b 🖌	
	Type of patients	

Figure 2: Bar diagram showing distribution of patients as per epileptic state.

Duration of illness	Number of patients	Percentage
<12 month	1013	21.7
1-2 yrs	1662	35.7
2-3 yrs	804	17.2
3-4 yrs	672	14.4
4-5 yrs	120	2.7
>5 yrs	381	8.3

Table 3: Distribution of patients as per duration of illness.

Therapeutic regime	Number of patients	Percentage
Phenobarbitone+Carbamezipine	49	15.75
Phenytoin sodium+phenobarbitone	52	16.72
Sodium valproate+Phenobatbitone	57	18.32
Sodium valproate+Phenytoin sodium	73	23.47
Sodium valproate+Carbamezipine	43	13.82
Phenobarbitone+Phenytoin sodium & Carbamezipine	37	11.9
	311	

Table 4: Distribution of old resistant patients as per their therapeutic regime.

Presenting feature	Number of patients	Percentage
Convulsion with unconsciousness frothing from mouth and autonomic dysfunction	607	13.05
Convulsion with unconsciousness and Frothing from mouth	2002	42.97
Convulsion with transient unconsciousness	1394	29.93
Mental debility	970	20.82
Physical debility	970	20.82

Table 5: Distribution of patients as per clinical presentation.

 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %
 %</t

and the production of the prod

مندر الارالي مرزان الس	%,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	%, /
in property and the second sec	····,···· %,··, * . , *	%

Page 2 of 4

Particulars	Number of patients	Percentage
Hematological:		
Hemoglobin-(%)		
8-10 gm	1957	42.01
10-12 gm	2423	52.01
12-14 gm	279	5.98
Hepatic parameters:		
Serum bilirubin-		
> 1 mg%	49	1.05
<1 mg%	4609	98.95
Renal parameters:		
Blood urea-		
>30 mg%	58	1.24
<30 mg%	4600	98.76
Serum creatinine		
>1.5 mg%	54	1.16
<1.5 mg%	4604	98.84

 Table 6: Distribution of patients as per their base hemato, hepatic and renal parameters.

, e company, e

Di c .. i

and a construction of the second construction and the second construction of the second construction o

Page 3 of 4

and a state of the second s

م از سام روید استان از در این به روید از ۲۰ میلی () این میتران از صورت از روید از میتران با میڈر میلی ۲۰۱۰ از این میتران به این روید این میتران در میتران در میتران در م ۲۰ مامری به در این میتران از در در میتران از میتران ۲۰ میتران در میتران از میتران از میتران در میتران از میتران از میتران از میتران از میتران از میتران از میتران

•••• ••• •••

And the party sugart.

an analog and the second and the second and the second and the second second second second second second second And the second second second second to second se And the second second

-1

References

- 1. Shankar A, Shubham (2010) Epidemiological study of epilepsy among children in Jharkhand. The Holistic pediatrics 4: 230-236.
- 2. Shankar A (2013) Ayurveda for neurological disorder. J Homeop Ayurv Med 2: 130-131.

Page 4 of 4

- Shankar AC, Shankar AA, Shankar SA (2008) Epileptics debility a clinician idiosyncrasy. IJM Today 4: 143-145.
- 4. Shankar A (2009) Analysis of clinicopathology of grand mal epilepsy and present trend of its management. India epilepsy forum bulletin 4: 212-216.
- Andrews DT, Sconfeld WH (1992) Predictive facts for controlling seizure using a behavioural approach. Seizure 1:111-116.
- Shankar A (2013) Chemical constituents and pharmacodynamics of herbal constituents: The Pharmacological basis of Indigenous therapeutics. Bhalani Publication, Mumbai, India.