

Bacterial Lipopolysaccharide-Dependent Complement Evasion

Bożena Futoma-Koloch*

Department of Microbiology, University of Wrocław, Przybyszewskiego Wrocław, Poland

*Corresponding author: Bożena Futoma-Koloch, Department of Microbiology, University of Wrocław, Przybyszewskiego Wrocław, Poland, Tel: 48 71 375 62 22; Fax: 48 71 325 21 51; E-mail: bozena.futoma-koloch@uw.edu.pl

Received date: May 01, 2016; Accepted date: May 02, 2016; Published date: May 07, 2016

Copyright: © 2016 Futoma-Koloch B, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Editorial

Finding factors that control complement (C) components deposition on bacterial cells is the way to explain the sophisticated mechanism of bacteraemia. Lipopolysaccharide (LPS, endotoxin) is a surface antigen found in the cell walls of Gram-negative bacteria, such as *Salmonella* [1]. LPS action on the human body is often considered in the context of sepsis, which is characterized by an excessive inflammatory response in the presence of endotoxin. At low concentrations, LPS stimulates innate immune system, which helps the host to eliminate the invader; but at high levels, it causes shock and even death [2]. Moreover, recent LPS characterization supports the development of a new vaccine against non-typhoidal *Salmonellae* for Africa [3]. From the point of view of pathogens, LPS-dependent resistance to C-mediated killing is an essential virulence property. In this paper, I want to highlight the main findings on the role of LPS in bacterial C-dependent serum susceptibility. To introduce, it is worth bringing forward a general structure of endotoxin. LPS of smooth strains (S) consists of three structural domains: lipid A, which anchors the whole molecule in the bacterial cell wall, core oligosaccharide, and O-specific polysaccharide chains (O antigen, O-Ag). Lipooligosaccharide (LOS) is analogous to the LPS, but it lacks O-Ag. When the LPS is connected to the bacterium, the outermost fragment - O antigen determines the response of the inflammatory system. In case of killing or disruption of bacterial cells with drugs (i.e. antibiotics) results in a lipid A release, the most toxic domain of endotoxin [1]. The role of LPS and its chemotypes in the resistance of bacteria to C has not been investigated thoroughly. Since the 80s, it has been known that long-chain LPS with complete O-specific chains confers bacterial resistance to serum by promoting the deposition of C components at a distance from the cell wall, thus preventing its disruption with the C5b-9 complexes. It was demonstrated that the amount of LPS O-Ag and its chain length distribution are important factors that protect f

t a s cer] ~ e
ø t,h] o b t h d th

7. Nelson BW, Roantree RJ (1967) Analyses of lipopolysaccharides extracted from penicillin-resistant, serum-sensitive salmonella mutants. *J Gen Microbiol* 48: 179-188.
8. Kaca W, Literacka E, Sjöholm AG, Weintraub A (2000) Complement activation by *Proteus mirabilis* negatively charged lipopolysaccharides. *J Endotoxin Res* 6: 223-234.
9. Cooper NR (1983) Activation and regulation of the first complement component. *Fed Proc* 42: 134-138.
10. Bjornson AB, Bjornson HS (1977) Activation of complement by opportunist pathogens and chemotypes of *Salmonella minnesota*. *Infect Immun* 16: 748-753.
11. Devyatyarova-Johnson M, Rees IH, Robertson BD, Turner MW, Klein NJ, et al. (2000) Lipopolysaccharide structures of *Salmonella enterica* serovar Typhimurium and *Neisseria gonorrhoeae* determine the attachment of human mannose-binding lectin to intact organisms. *Infect Immun* 68: 3894-3899.
12. Biedzka-Sarek M, Venho R, Skurnik M (2005) Role of YadA, Ail, and lipopolysaccharide in serum resistance of *Yersinia enterocolitica* serotype O:3. *Infect Immun* 73: 2232-2244.
13. Grossman N, Leive L (1984) Complement activation via the alternative pathway by purified *Salmonella* lipopolysaccharide is affected by its structure but not its O-antigen length. *J Immunol* 132: 376-385.
14. Fudala R, Doroszkiewicz W, Niedbach J, Gamian A, Weintraub A, et al. (2003) The factor C3 conversion in human complement by smooth *Shigella flexneri* lipopolysaccharides. *Acta Microbiol Pol* 52: 45-52.
15. Lewis LA, Gulati S, Burrowes E, Zheng B, Ram S, et al. (2015) -2,3-Sialyltransferase expression level impacts the kinetics of lipooligosaccharide sialylation, complement resistance, and the ability of *Neisseria gonorrhoeae* to colonize the murine genital tract. *mBio* 6: e02465-14.
16. Ram S, Sharma AK, Simpson SD, Gulati S, McQuillen DP, et al. (1998) A novel sialic acid binding site on factor H mediates serum resistance of sialylated *Neisseria gonorrhoeae*. *J Exp Med* 187: 743-752.
17. Bugla-Plosko ska G, Rybka J, Futoma-Koloch B, Cisowska A, Gamian A, et al. (2010) Sialic acid-containing lipopolysaccharides of *Salmonella* O48 strains-potential role in camouflage and susceptibility to the bactericidal effect of normal human serum. *Microb Ecol* 59: 601-613.
18. Futoma-Koloch B, Godlewska U, Guz-Regner K, Dorotkiewicz-Jach A, Klaus E, et al. (2015) Presumable role of outer membrane proteins of *Salmonella* containing sialylated lipopolysaccharides serovar Ngozi, sv. Isaszeg and subspecies arizonae in determining susceptibility to human serum. *Gut Pathog* 7: 18.
19. Ellström P, Feodorof B, Hänninen M-L, Rautelin H (2014) Lipooligosaccharide locus class of *Campylobacter jejuni*: sialylation is not needed for invasive infection. *Clin Microbiol Infect* 20: 524-529.