

# Beta Cells in Pancreas

Raveendran AV E

Department of General Medicine, University of Calicut, Calicut, India

\*Corresponding author:

Received date: 10/10/2023 Accepted date: 10/10/2023 Published date: 10/10/2023

Citation: Raveendran AV E. Beta Cells in Pancreas. *Journal of Health and Medical Research*. 2023;10(10):1000-1005.

Copyright: © 2023 Raveendran AV E. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Editorial

Beta cells (β-cells) are a type of cell found in pancreatic islets that synthesize and secrete insulin and amylin. Beta cells make up 50-70% of the cells in human islets. In patients with type 1 diabetes, beta-cell mass and function are diminished, leading to insufficient insulin secretion and hyperglycemia. The primary function of a beta cell is to produce and release insulin and amylin. Both are hormones which reduce blood glucose levels by different mechanisms [1]. Beta cells can respond quickly to spikes in blood glucose concentrations by secreting some of their stored insulin and amylin while simultaneously producing more. Beta cells are the only site of insulin synthesis in mammals. As glucose stimulates insulin secretion, it simultaneously increases proinsulin biosynthesis, mainly through translational control. The insulin gene is first transcribed into mRNA and translated into preproinsulin. After translation, the preproinsulin precursor contains an N-terminal signal peptide that allows translocation into the rough endoplasmic reticulum. Inside the RER, the signal peptide is cleaved to form proinsulin. Then, folding of proinsulin occurs forming three