of paramount importance [7,8]. Bioremediation can be de ned as any process that uses microorganisms (Bacteria,fungi), green plants or their enzymes to return the natural environment altered bycontaminants of original condition. Bioremediation constitutes an attractive alternative to physico-chemical methods of remediation, as it is less expensive and can selectively achieve complete destruction of organic pollutants [9]. Microorganisms are thought to play an important role in the removal and detoxi cation of these toxicants from the environment. Degradation of pesticides is usually bene cial, since the reactions that destroy pesticides, convert most pesticide residues in the environment to inactive, less toxic, harmless compounds [10].

For many degradation studies Pseudomonas sp. was selected as it is a well-known biodegrader of agrochemicals [11]. Five bacterial isolates (Pseudomonas aeruginosa, Klebsiella Sp., Escherichia coli, Bacillus S and Corynebacterium) isolated from brinjal cultivated eld were found

Keywords Bioremediation; Methomyl; Pseudomonas aeruginosa; Plasmid

Introduction

A wide variety of synthetic chemicals have found their way into the ecosystem as a consequence of industrial activities, agricultural applications and use in domestic purpose [1]. Pesticides are organic compounds manufactured and used for pest control. When pesticides are dispersed in the environment, they become pollutants, with ecological e ects that require remediation. Environmental pollution is caused by both excessive and continuous use of pesticides,xcessmÿ5rfs w0hurtfound their way into cause contamination of both ground and surface water resources [6]. e residual pesticide comes in contact with water causing surface/ ground water pollution leading to the toxicity of biotic environment. erefore, bioremediation techniques for the treatment of pesticides are

methomyl further (Table 5) which indicates that the degradative property of methomyl was encoded in the plasmid of Pseudomonas aeruginosa.

Discussion

e discovery of microorganisms capable of tolerating or growing in high concentrations of pesticides provides a potentially interesting avenue for treating hazardous wastes [4]. Some investigations resulting in the identi cation of microbial isolates which are apparently responsible for the accelerated degradation of individual pesticides is necessary [30,31] of which the 16s rDNA analysis is considered to the most valid and widely used technique in the identi cation of microorganisms [8,32-34].

In the present study, the most potential bacterial strain with the highest ability to degrade methomyl was identi ed morphologically, biochemically and genetically. ey were gram negative, rod shaped bacteria, they were Indole, MR and VP negative, H_2S negative, citrate, catalase, gelatin hydrolysis and oxidase tests were positive and the analysis of the 16S rDNA showed highest similarity to Pseudomonas aeruginosa.

Biodegradation of methomyl by *Pseudomonas aeruginosa*

e present study revealed that the methomyl was decreased signi cantly by Pseudomonas aeruginosa. However, di erent degradation rates were observed. All the organisms could utilize methomyl as a sole carbon source and had the bioremediation ability. Further it was observed that the soil isolate Pseudomonas aeruginosa, exhibited a good growth rate and higher biodegradation e ciency towards methomyl when compared to the other isolates demonstrating their potential use in bioremediation. It is suggested that the detoxi cation metabolism occurs when a microorganism uses the pesticide as a carbon and energy source and the process facilitated by resistant microorganisms [34]. Earlier studies have reported that Pseudomonas is a versatile genus and suggested that th genus could degrade a number of chemicals like pesticides including carbaryl, malathion, p-nitrophenol and parathion, bethoxazin, it is widely present in soil and can be used to clean up di erent xenobiotics compounds like propiconazole [35]. Earlier, two microorganisms classi ed as Pseudomonas sp. [36] and Escherichia coli [37] have been isolated and described as possessing the ability to degrade lindane. e results obtained in this study were in agreement with earlier reports that indicated the inVolvement of di erent species of Enterobacteriaecae in the degradation of insecticides like chlorpyrifos, phosphonate and glyphosate [38]. Several Pseudomonas sp. that metabolize organophosphorous and carbamate pesticides have been isolated from soil [30,31]. It has been reported that Stenotrophomonas maltophilia M1 strain is capable of methomyl degradation [39]. Bhagobaty and Malik [40] have reported of the Pseudomonas strains capable of degrading chlorpyrifos. Bioremediation of endosulfan was carried out by Staphylococcus sp and

47. 6HUGDU *LEVRQ '7 0XQQHFNH '0 /DQFDVWHU 48 6LQJK 3DDNHU $ 0RUJDQ -$:ULJKW '- (IIHFW ,QYROYHPHQW LQ 3DUDWKLRQ +\GURO\VLV E\ 3VHXGRPRQDV GLPLQXWD $SSO (QYLURQ Microbiol 44: 246-249.