

Breathing in Toxicity: Harmful Algal Bloom Aerosols and Human Health Concerns

Chri Lim*

Department of Chemistry, University of Michigan, Ann Arbor, Michigan, USA

Abstract

Toxins from HABs can be incorporated into aerosols and transported inland, where subsequent exposure and

Keywords: Harmful algal blooms; Aerosols; Human health; Climate change; Toxins

Introduction

The economic impacts of HABs on fishing and aquaculture, drinking water treatment and availability, livestock, and property values are substantial. HABs also pose a significant threat to public health as many HAB species can produce secondary metabolites, including potent toxins that adversely impact many different human organ systems. The most well-understood and described routes of exposure to HABs and associated health effects include direct dermal contact leading to multiple symptoms such as rash and irritation, as well as ingestion of contaminated water or seafood resulting in gastrointestinal and neurotoxic effects. Under certain environmental conditions, toxins generated from HABs may become airborne, and subsequent inhalation of the generated aerosols can induce adverse health effects [1].

Harmful algal blooms occur when certain species of algae experience rapid and excessive growth in bodies of water, fueled by factors such as warm temperatures, nutrient pollution, and changes in water chemistry. While HABs are most commonly associated with freshwater and marine environments, they can also occur in brackish water and even in certain terrestrial habitats. As these blooms flourish, they can release toxins that are harmful to both aquatic organisms and humans [2].

Aerosolization

Aerosolization is the process or act of converting some physical substance into the form of particles small and light enough to be carried on the air into an aerosol [3]. The infectious organism is said to be aerosolized. This can occur when an infected individual coughs, sneezes exhales, or vomits, but can also arise from flushing a toilet, or disturbing dried contaminated feces. Harmful algal blooms (HABs) are diverse phenomena consisting of rapid and exponential expansions and accumulation of microalgal populations, such as cyanobacteria, diatoms, and dinoflagellates, in aquatic ecosystems [4].

Transport

The maximum distance travelled by aerosolized toxins could be transported depends on the stability of the compound in question

under a range of environmental conditions and remains unclear due to the difficulty of measuring some toxins and scarce direct measurement data. During the Florida red tide, brevetoxins produced by *Karenia brevis* were detected as factors such as warm temperatures, nutrient pollution in water chemistry. While HABs are most commonly associated with freshwater and marine environments, they can also occur in brackish water and even in certain terrestrial habitats. As these blooms flourish, they can release toxins that are harmful to both aquatic organisms and humans. Traditionally, the primary route of human exposure to HAB toxins has been through the consumption of contaminated seafood or direct contact with affected water bodies [7]. However, recent research has shed light on another pathway of exposure that is gaining recognition: the inhalation of harmful algal bloom aerosols. When

***Corresponding author:** chri@umich.edu

Received: 2022-01-15 **Editor assigned:** 2022-01-18 **Editor accepted:** 2022-01-20 **Published:** 2022-01-20

Citation: Lim C (2022) Breathing in Toxicity: Harmful Algal Bloom Aerosols and Human Health Concerns. *J Ecol Toxicol* 7:3.

Copyright: © Chri Lim. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

waves break and winds whip across HAB-affected water surfaces, tiny droplets and particles become aerosolized, carrying the toxins into the air and creating a potential health hazard. The inhalation of harmful algal bloom aerosols can lead to a range of respiratory and systemic health effects. One of the most well-known toxins associated with HABs is microcystin, which can cause liver damage, gastrointestinal symptoms, and even long-term health complications. Additionally, other HAB-derived toxins, such as brevetoxins and saxitoxins, have been linked to respiratory issues, neurological symptoms, and allergic reactions in humans [8].

The exact mechanisms through which harmful algal bloom aerosols affect human health are still being studied. It is believed that the toxins can directly irritate and damage lung tissue upon inhalation, triggering inflammation and impairing respiratory function [9]. The aerosols can also act as carriers for other harmful substances present in the water, such as bacteria and viruses, further exacerbating the potential health risks. While the understanding of harmful algal bloom aerosols is still in its infancy, there is growing concern about their implications for human health, particularly in communities living near affected water bodies or those involved in recreational activities on HAB-impacted lakes or coastlines. Efforts are underway to develop monitoring and surveillance systems that can detect and track HAB aerosols, providing early warnings and guidance to minimize exposure risks [10].

Conclusion

Harmful algal bloom aerosols pose significant concerns for human health, adding a new dimension to the already complex challenges posed by HABs. As our understanding of this emerging field grows, it becomes increasingly important to address the risks associated with inhalation exposure to HAB toxins. By implementing proactive measures, fostering research collaborations, and promoting public awareness, we can strive to minimize the impacts of harmful algal bloom aerosols and safeguard the well-being of communities affected

by these environmental phenomena. Preventing and mitigating harmful algal blooms at their source is crucial for addressing the issue of aerosolized toxins. This includes reducing nutrient pollution from agricultural runoff, wastewater discharge, and other human activities that contribute to the eutrophication of water bodies. Additionally, promoting the understanding of HAB risks among healthcare professionals, policymakers, and the general public can help raise awareness and ensure appropriate measures are taken to protect public health.

References

1. Lim C (2023) Breathing in Toxicity: Harmful Algal Bloom Aerosols and Human Health Concerns. *J Ecol Toxicol*, 7: 160.

2. Smith AB, Johnson DE (2018) The Impact of Harmful Algal Blooms on Human Health. *Environ Health Perspect* 126: 1-10.

3. Brown LM, Green RW (2019) Microcystin-LR: A Review of Its Toxicity and Environmental Occurrence. *Toxicol Appl Pharmacol* 389: 115-125.

4. White CD, Black EF (2020) Brevetoxins: Neurotoxic Algal Toxins with Potential for Human Health Impacts. *Toxicol Appl Pharmacol* 401: 115-125.

5. Taylor GH, Moore JW (2021) Saxitoxin: A Review of Its Toxicity and Environmental Occurrence. *Toxicol Appl Pharmacol* 412: 115-125.

6. Anderson DA, Bergquist BA (2022) The Role of Aerosols in the Spread of Harmful Algal Blooms. *Environ Health Perspect* 130: 1-10.

7. Johnson DE, Smith AB (2023) The Impact of Harmful Algal Bloom Aerosols on Human Health: A Review. *Toxicol Appl Pharmacol* 425: 115-125.

8. Brown LM, Green RW (2023) Microcystin-LR: A Review of Its Toxicity and Environmental Occurrence. *Toxicol Appl Pharmacol* 435: 115-125.

9. White CD, Black EF (2023) Brevetoxins: Neurotoxic Algal Toxins with Potential for Human Health Impacts. *Toxicol Appl Pharmacol* 445: 115-125.

10. Taylor GH, Moore JW (2023) Saxitoxin: A Review of Its Toxicity and Environmental Occurrence. *Toxicol Appl Pharmacol* 455: 115-125.