Carrier Frequency of Congenital Leptin Deficiency in Central Punjab Region of Pakistan

Muhammad Wasim* **and Nida Fakhar**

Institute of Biochemistry and Biotechnology, University of Veterinary and Animal Sciences Lahore, Pakistan

***Corresponding author:** Muhammad Wasim, Institute of Biochemistry and Biotechnology, University of Veterinary and Animal Sciences, Lahore, Pakistan, Tel: 923224990977; E-mail: mm.waseemjee@gmail.com

Received date: March 24, 2015; **Accepted date:** April 25, 2015; **Published date:** April 30, 2015

Copyright: © 2015 Wasim M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Congenital leptin deficiency is a rare autosomal recessive disorder, characterized by hyperphagia and early onset obesity. Worldwide data on this disorder includes only 25 cases out of this only six pathogenic mutations have been reported in the leptin gene. Most of these cases have been reported in Central Punjab region of Pakistan, harboring a (p.Gly133fsX145) mutation in homozygous state. This mutation has founder effect in Arain caste from Central Punjab region of Pakistan. In this study, carrier frequency of (p.Gly133fsX145) mutation was determined in 50 individuals belonging to Arain caste. Analysis of amplicons contains p.Gly133fsX145 mutation were analyzed by different techniques such as SSCP, Heteroduplex analysis and sequencing. Mutation was not found in any individual studied. This finding suggests that frequency of (p.Gly133fsX145) mutation in Arain caste is not as higher as its predicted founder effect in congenitally obese children of this caste was anticipated.

Keywords: Congenital leptin deficiency; Hyperphagia; SSCP; Heteroduplex analysis; p.Gly133fsX145

Introduction

Leptin is a 16-kDa adipocytokine or adipokine hormone which has 167 amino acids [1]. Firstly discovered in mice in 1994, the gene is known as Ob and the mutated mice known as Ob/Ob mice [2]. Mutation in leptin gene of Ob/Ob mice was first explained in the 1950s, a morbidly obese phenotype is caused by spontaneous mutation due to more eating habit and less energy expenditure [3].

Circulating leptin level decrease during fasting and increase during feeding, which indicates that leptin is a physiological signal for transition between the states of energy adequacy and starvation [4,5]. Role of this hormonal protein is like a satiety factor and it helps in regulating energy balance. It regulates the reproductive, hematopoietic, and immune systems. It is involved in development of brain and modulation of carbohydrate metabolism [6]. The most important pathway is leptin-melanocortin pathway presently known to be involved in five congenital forms of monogenic obesity [7].

Leptin is a neurotransmitter which acts via two populations of arcuate (Arc) neurons located in the hypothalamus. The first group of neurons expresses Agouti-related Peptide (AgRP) and Neuropeptide Y (NPY) and the second one expresses Pro-Opio-Melano-Cortin (POMC) and Cocaine and Amphetamine-Related Transcript (CART). Leptin acts on POMC/CART neurons to signal repletion of energy stores and thereby causes suppression of food intake and increase in energy expenditure. The action of leptin on AgRP/NPY neurons is inhibitory and their activation through fasting increases food intake and energy conservation [7,8]. Congenital leptin deficiency is caused by mutations in the leptin gene that cause obesity. Until now only 6 pathogenic leptin mutations have been reported.

The first homozygous mutation was (p.Gly133fsX145) detected in the leptin gene of two Pakistani cousins who had undetectable serum leptin level [9]. The second leptin mutation (R105W) was identified in 4 members of a Turkish family (Strobel et al. 1998), the third (N103K) in 2 Egyptian children [10] and the fourth (L72S) in an Austrian 14 years old female child [11]. Fifth and sixth mutations were (p.Leu161fsX170 and c.104_106delTCA) in obese children of Pakistan [12]. Congenital leptin deficiency has a pleiotropic effect and also associated with multiple disorders such as deregulation of reproductive, immune, neuroendocrine and metabolic physiology [9,13-15].

Most of the reported cases of congenital leptin deficiency had Pakistani origin; the disorder might be a frequent cause of morbid childhood obesity in Pakistan, based on higher rates of consanguineous marriages in Central Punjab [12]. Except two of these cases, all harbored (p.Gly133fsX145) mutation in Arain caste indicating founder effect. Therefore the present study was designed to