Chronic Pain Management and Its Relationship to Physiological Variables

Eleni G Hapidou[1*] **and Kayli M Culig**[2]

[1]*Michael G DeGroote Pain Clinic, Hamilton Health Sciences & Michael G DeGroote Institute for Pain Research and Care, & Department of Psychiatry and Behavioral Neurosciences, & Department of Psychology, Neuroscience and Behavior, McMaster University, 1200 Main St. West, Hamilton, ON L8N 3Z5, Canada*

[2]*Bachelor of Health Sciences (Honor's) Program, McMaster University, 1280 Main Street West, Hamilton, Ontario L8S 4L8, Canada*

***Corresponding author:** Eleni G Hapidou, Michael G. DeGroote Pain Clinic, McMaster University Medical Centre, 4th Floor, Yellow Section 4V, 1200 Main St. West, Hamilton, ON L8N 3Z5, Canada, Tel: (905) 521-2100, Ext. 77492; Fax : (905) 577-8022; E-mail: hapidou@hhsc.ca

Received date: January 27, 2019; **Accepted date**: February 16, 2019; **Published date:** February 23, 2019

Copyright: © 2019 Hapidou EG. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Chronic pain (CP) is defined as pain lasting more than 3 months. It affects thousands of Canadians daily through biological, psychological and social factors. Not only are physiological factors affected in those who experience chronic pain but also sleep, mood, and general quality of life. We do not yet know the exact biological mechanisms through which acute pain and injury develop into chronic pain, however, in this article; we discuss a dominant hypothesis that might offer an explanation: Central Sensitization. In addition, the purpose of this article is to explore the biological mechanisms of chronic pain and the importance of using physiological measures to assess the outcomes of pain management programs. This paper draws attention to the importance of having further research conducted in order to understand the underlying biological causes of chronic pain as well as identifying specific biomarkers that can be used to measure treatment outcomes. This will allow us to design effective and innovative pain management programs in order to improve the quality of life for CP patients.

Keywords: Chronic pain; Stress response; Central sensitization; Pain management; Physiological measures

b R b

R S R

Chronic pain (CP) is defned as pain lasting ≥3 months and its subjective perception is complex: it is comprised of interacting biological, psychological and social factors [1]. CP continues to be an ongoing challenge in Canada, afecting as many as 20% to 29% individuals nationwide [2]. For those living with a CP condition, it can provoke signifcant long-term debilitation and sufering [3]. In particular, CP can negatively afect many domains of a patient's health including sleep, cardiovascular ftness, mood, sexual functioning and overall quality of life [1]. Among Canadians waiting for efective intervention to relieve their CP, over two thirds reported 'severe pain' (i.e., ≥7 out of 10 on a Likert scp p dis n og ' quality of life and daily functioning [4]. CP also poses enormous economic burden on individual and societal levels. For example, Canadians with CP awaiting treatment reported an average median monthly cost of $1,462 (CDN) for care [5]. In a newly publicized population-based study [6], they found that the incremental healthcare costs amounted to 50% higher in patients managing CP than their healthy control counterparts. From a broader perspective, Canada spends approximately $6 billion annually on direct CP expenditure, and $37 billion annually on indirect costs (i.e., loss of job productivity, loss of jobs, employee sick days etc.) [7]. Evidently, more resources and research should be directed at combatting this pressing health concern.

T e a b s a a s a a s

It has been established that CP may develop as a result of a dysfunctional stress response [1]. Normal functioning, interdependent systems (i.e., nervous, endocrine and immune) interact to adaptively respond to an acute stressor or injury. T is bodily response is known as allostasis and is necessary to maintain homeostasis, thereby protecting vital internal processes. When presented with a stressor, allostatic systems such as the hypothalamic-pituitary-adrenal (HPA) axis and autonomic nervous system (ANS) promote a 'fght or fight' response so that individuals can efectively respond to the homeostatic imbalance. In many cases, the HPA axis facilitates the release of

waitlists of multidisciplinary pain treatment facilities? Can J Anaesth 57: 539- 548.

5. Guerriere DN, Choinière M, Dion D, Peng P, Staford-Coyte E, et al. (2010) The Canadian STOP-PAIN project-Part 2: What is the cost of pain for patients on waitlists of multidisciplinary pain treatment facilities? Can J Anaesth 57: 549- 558.
6. Hogan ME, Taddio A, Katz J, Shah V, Krahn M (2016) Incremental health care costs for chronic pain in Ontario, Canada: A population-based matched cohort study of adolescents and adults using administrative data. Pain 157: 1626-1633.
7. Lynch ME (2011) The need for a Canadian pain strategy. Pain Res Manag 16: 77- 80.
8. Gupta A, Silman AJ (2004) Psychological stress and fbromyalgia: a review of the evidence suggesting a neuroendocrine link. Arthritis Res Ther, 6: 98.
9. Selye H (1956) The stress of life. McGraw-Hill Book Company. J Bone joint Surg 39: 479.
10. McEwen BS (1998) Stress, adaptation, and disease: Allostasis and allostatic load. Ann N Y Acad Sci 840: 33-44.
11. Hannibal KE, Bishop MD (2014) Chronic stress, cortisol dysfunction, and pain: a psychoneuroendocrine rationale for stress management in pain rehabilitation. Phys Ther 94: 1816.
12. Chapman CR, Tuckett RP, Song CW (2008) Pain and stress in a systems perspective. J Pain 9: 122-145.
13. Mifin KA, Kerr BJ (2014) The transition from acute to chronic pain: understanding how diferent biological systems interact. Can J Anaesth 61: 112-122.
14. Adler GK, Geenen R (2005) Hypothalamic- pituitary- adrenal and autonomic nervous system functioning in fbromyalgia. Rheum Dis Clin North Am 31: 187-202.
15. Raison VMCL (2009) Neurobiology of depression, fbromyalgia and neuropathic pain. Front Biosci 14: 5291-5338.
16. Wolfe F, Smythe HA, Yunus MB, Bennett RM, Bombardier C, et al (1990) The American College of Rheumatology 1990 criteria for the classifcation of fbromyalgia. Arthritis Rheum 33: 160-172.
17. Lee KE, Choi SE, Kang JH, Yim YR, Kim, JE, et al. (2016) Comparison of heart rate variability and classic autonomic testing for detection of cardiac autonomic dysfunction in patients with fbromyalgia. Int J Rheum Dis 21: 804-812.
18. Martínez-Lavín M, Hermosillo AG, Rosas M, Soto ME (1998) Circadian studies of autonomic nervous balance in patients with fbromyalgia: a heart rate variability analysis. Arthritis and Rheumatism 41: 1966-1971.
19. Addison RG (1984) Chronic pain syndrome. Americ J Med 77: 54-58.
20. Hainline B (2005) Chronic pain: physiological, diagnostic, and management considerations. Psychiatric Clinics of North America, 28: 713-735.
21. Vanegas H, Schaible HG (2004) Descending control of persistent pain: Inhibitory or facilitatory? Brain Res Brain Res Rev 46: 295-309.
22. Dubner R (2004) The neurobiology of persistent pain and its clinical implications. Clin Neurophysiol 57: 3-7.
23. Herman JP, Figueiredo H, Mueller NK, Ulrich-Lai Y, Ostrander MM, et al. (2003) Central mechanisms of stress integration: Hierarchical oO In IarP o, co tag r eEyp o n v nsg u o N eur o en