

INTRODUCTION

Panic disorder is one type of anxiety disorder of anxiety disorders according to the Diagnostic and Statistical Manual of

“Antisocial personality disorder”.

NEO-PI-R

Incorporated Medical Institution Nagoya Mental Clinic, and the members of the Department of Psychiatry, Division of Neuroscience, Graduate School of Medicine, Mie University. We wish to thank Yukika Nishimura, PhD (previously, Department of Psychiatry, Division of Neuroscience, Graduate School of Medicine, Mie University, Mie, Japan) for valuable comments.

REFERENCES

- Bomyea, J., Lang, A.J., Craske, M.G., Chavira, D., Sherbourne, C.D., Rose, R.D., et al. (2013). Suicidal ideation and risk factors in primary care patients with anxiety disorders. , 60-65.
- Cabinet, Offce. (2015). [Heisei27nenbanjisatsutsaisakuhauso]. Shobi Printing Company Ltd, 219 [in Japanese].
- Diaconu, G. & Turecki, G. (2007). Panic disorder and suicidality: is comorbidity with depression the key? , 203-209.
- Hara, N., Nishimura, Y., Yokoyama, C., Inoue, K., Nishida, A., Tanii, H., et al. (2012). The development of agoraphobia is associated with the symptoms and location of a patient's frst panic attack. , 12.
- Hirata, M., Kawanishi, C., Oyama, N., Miyake, Y., Otsuka, K., Yamada, T., et al. (2013). Training workshop on caring for suicide attempters implemented by the Ministry of Health, Labour and Welfare, Japan. : 64.
- Iida, T., Inoue, K., Ito, Y., Ishikawa, H., Teradaira, R., Ota, A., et al. (2011). [Kokkasikensutoresunesunioruyokuutsu fuankeikounokoudoutokusei (Dai ippou)]. , 13-14. [in Japanese]
- Inoue, K. & Fujita, Y. [Eds. Xenitidis, K., & Campbell, C.]. (2013). Effective suicide prevention measures for teenagers in Japan. , 312.
- Inoue, K., Fukunaga, T., Nata, M., Abe, S., Okazaki, Y. (2013). Discussion of extensive suicide prevention based on Suicide statistics from 2006 to 2009 in Mie Prefecture, Japan. , 646-648.
- Inoue, K., Tanii, H., Nishimura, Y., Masaki, M., Yokoyama, C., Kajiki, N., et al (2007). Suicide in panic disorder. , 199-202.
- Nishimura, Y., Tanii, H., Hara, N., Inoue, K., Kaiya, H., Nishida, A., et al. (2009). Relationship between the prefrontal function during a cognitive task and the severity of the symptoms in patients with panic disorder: a multi-channel NIRS study. , 168-172.
- Ohtani, T., Kaiya, H., Utsumi, T., Inoue, K., Kato, N., Sasaki, T. (2006). Sensitivity to seasonal changes in panic disorder patients. , 379-383.
- Otsubo, T. (2016). Brief structured interview for psychiatric disorders; M.I.N.I. and M.I.N.I. KID. , 16-21 [in Japanese].
- Shimonaka, Y., Nakazato, K., Gono, Y., Takayama, M. (1998). Construction and factorial validity of the Japanese NEO-PI-R. , 138-147 [in Japanese].
- Takeshima, T., Yamauchi, T., Inagaki, M., Kodaka, M., Matsumoto, T., Kawano, K., et al. (2015). Suicide prevention strategies in Japan: a 15-year review (1998-2013). , 52-66.