Data collection

We measured the anthropometric parameters of each participating child, such as weight and height, following standard protocols. For the measurement of weight, the children were advised to remove any heavy clothing and shoes whereas for the measurement of height they were instructed to stand in the right position. The measurement of height and weight was performed at the moment the patient showed up at our hospital (that is, before the surgical treatment) and then 6 months and 12 months after the surgical treatment of chronic tonsillitis. The same calibrated weight and height scales were used to measure the weight and height of children in each of the three measurement points of time. Also, we used the standard reference levels of weight and height by age, in order to compare our data against such thresholds.

Data analysis

For categorical variables absolute numbers and the respective percentages were reported. For continuous variables the mean value and the standard deviation of the parameter was reported. To compare the mean values of weight and height in different points in time the student's t-test, through the one-way ANOVA procedure, was used. For categorical variables, the chi-square test was used. The associations were considered as statistically significant if the P-value ≤0.05.

All the statistical analysis were carried out using the Statistical Package for Social Science (SPSS) software, version 16.

Results

There were 85 children that participated in the study, of whom 37 were boys (43.5%) and 48 girls (56.5%). The mean age of participants was 7.15 years ± 1.89 years, without significant difference between boys and girls (P=0.849) (Table 1). The age of the participants ranged from 4 years (8 subject

Variable	Total		Boys (n=37)		Girls (n=48)	
	Mean value ± SD	**P- value**	**Mean value ± SD**	**P- value**	**Mean value ± SD**	**P- value**
Weight (in kg)				0.367		0.538
Before the operation	23.86 ± 7.33 24.96 ± 7.39	0.212*	24.27 ± 7.03		23.54 ± 7.61	
6 months after operation	25.87 ± 7.51		25.53 ± 6.99		24.52 ± 7.73	
12 months after operation			26.61 ± 7.20		25.29 ± 7.77	
Height (in cm)		0.012		0.112		0.105
Before the operation	119.9 ± 11.2		120.3 ± 10.6		119.8 ± 11.8	
6 months after operation	122.6 ± 11.5		123.0 ± 11.0		122.4 ± 12.1	
12 months after operation	125.3 ± 11.8		125.8 ± 11.9		125.0 ± 11.9	

*P - value according to one-way ANOVA test.

Total	32.93	23.54	24.52	25.29	0.538	0.267
*P - value according to one-way ANOVA test.						

Table 3: Mean weight of the subject in the study according to age and gender and time of measurement.

Boys

Variable	**Standard mean height**	**Mean height before the operation**	**Mean height 6 months after the operation**	**Mean height 12 months after the operation**	**P-value for three periods**	**P- value before the op. 12 month after op.**
Age	109.5	102.0	103.9	106.4	0.164 *	0.106 *
4 years	116.8	110.6	112.9	114.7	0.020	0.018
5 years	123.9	116.7	118.2	119.6	0.074	0.031
6 years	131.0	119.2	123.0	125.4	0.017	0.013
7 years	137.8	124.2	128.2	132.5	0.005	0.002
8 years	144.1	130.8	132.8	135.0	0.515	0.287
9 years	149.9	134.6	137.6	141.2	0.176	0.087
10 years	155.5	135.0	140.0	149.0	-	-
11 years						
Total	131.2	120.3	123.0	125.8	0.112	0.032
Girls						
Age	108.4					
4 years	116.1					
5 years	123.9					
6 years	131.3					
7 years	137.7					
8 years	143.8					
9 years	14ţ					
10 years						

Height before operation Under the standard In norm	84 (98.8) * 1 (1.2)	37 (100.0) 0 (0.0)	47 (97.9) 1 (2.1)	0.377
Height 6 months after operation Under the standard In norm	84 (98.8) 1 (1.2)	37 (100.0) 0 (0.0)	47 (97.9) 1 (2.1)	0.377
Height 12 months after operation Under the standard In norm	81 (95.3) 4 (4.7)	36 (97.3) 1 (2.7)	45 (93.8) 3 (6.3)	0.444
*The absolute number and percentage according to columns (in parenthesis). †P - value according to hi square test				

Table 6: The patients' height in the study compared to standard reference levels.

Discussion

This is the first study addressing the association of chronic tonsillitis and anthropometric parameters such as weight and height among young children in Kosovo. We noticed that the mean weight and mean height of chronic tonsillitis children was lower than mean standard reference levels, for each age group and in both boys and girls. Also, we evidenced that the surgical treatment of chronic tonsillitis resulted in an improvement of mean weight and mean height of study participants and the improvement was greater about one year after the intervention, whereas for height these differences were significant in most cases. In addition, despite the lack of statistical significance regarding changes of weight, we spotted a clinical significance: the findings indicate that there is a positive association between time after surgical correction of chronic tonsillitis and weight and height of patients, suggesting the improvement of the