

Combining Epidemiology and Toxic genomics to Support an Unfocused Investigation of Pesticide Exposure and Parkinson's Disease

Madelline Garcia*

Toxic genomics; Pesticides; Parkinson's disease

For more than 50 years, pesticides have been used extensively in agriculture all around the world. At this time, the chemical industry had rapid growth, which facilitated the launch and widespread commercial use of numerous goods. For instance, 13,540 pesticide formulations and 1,074 various active chemicals are now approved for usage in the state of California (California Department of Pesticide Regulation, 2021). Modern commercial agriculture relies heavily on pesticides to assist maximise food output. However, because pesticides are deliberately made to kill living things (such as plants, fungus, insects, and rodents), they must be adequately evaluated for any possible negative effects on human health, especially when used on a large basis. The majority of the time, chronic exposures, mixtures, or sequential treatments with different pesticides are not taken into account in the toxicity testing used to register pesticides in the United States. Instead, it is usually based on single-product, single-target assessments using rodents as model organisms [1, 2].

three agricultural counties in Central California (Kern, Fresno, and Tulare) based on the study population's exposure prevalence [3,4].

chemical compounds that may interfere with bodily functions using in vitro biologic assays [7,8].

Several epidemiologic studies have shown that pesticides are among the environmental risk factors that are most consistently linked to Parkinson's disease (PD). Epidemiologic studies and experimental investigations, however, have not been able to fully evaluate long-term, low dose exposure related health effects for the majority of agents due to the needs of commercial agriculture for variety in agents that are tailored to emerging and reemerging pests as well as the large number of pesticides currently in use, i.e., 1074 active ingredients alone registered in California.

Overall, a range of specific pesticides have been linked to PD risk by our thorough, pesticide-wide association research and toxicologic/toxicogenomic integration. We were able to link pesticides to specific biologic and molecular targets relevant to the aetiology of Parkinson's disease (PD) thanks to cross-database queries we carried out and presented here. This information can help us better understand the involvement of pesticides in the genesis of Parkinson's disease (PD) and prioritise experimental pesticide research [9,10].

None.

None.

References

1. Reppel L, Schiavi J, Charif N, Leger L, Yu H, Pinzano A, et al. (2015) Chondrogenic induction of mesenchymal stromal/stem cells from Wharton's jelly embedded in alginate hydrogel and without added growth factor: an alternative stem cell source for cartilage tissue engineering. *Stem Cell Res Ther* 6: 260.

*Corresponding author: Madelline Garcia, Department of Animal Physiology, College of Colchester, United Kingdom, E-mail: Madelline33@gmail.com

Received: 03-Mar-2023, Manuscript No: tyoa-23-91159; **Editor assigned:** 06-Mar-2023, Pre-QC No: tyoa-23-91159 (PQ); **Reviewed:** 20-Mar-2023, QC No: tyoa-23-91159; **Revised:** 22-Mar-2023, Manuscript No: tyoa-23-91159 (R); **Published:** 29-Mar-2023, DOI: 10.4172/2476-2067.1000204

Citation: Garcia M (2023) Combining Epidemiology and Toxic genomics to Support an Unfocused Investigation of Pesticide Exposure and Parkinson's disease. *Toxicol Open Access* 9: 204.

Copyright:

A publicly accessible database called the CTD aims to advance our knowledge of how environmental exposures affect human health.

The database links information about chemicals, genes, phenotypes, diseases, and exposures to give contextualised knowledge about chemical exposures and human health through manual curation of peer-reviewed scientific literature. To keep the database complete and up to date, the CTD incorporates new research once a month [5, 6].

The Environmental Protection Agency (EPA), the National Toxicology Program (NTP) at NIEHS, the National Center for Advancing Translational Sciences (NCATS) at NIH, and the Food and Drug Administration (FDA) are working together as part of the Tox21 project to experimentally evaluate the toxicity of thousands of

2. Anzalone R, Lo Iacono M, Corrao S, Magno F, Loria T, et al. (2010) New emerging potentials for human Wharton's jelly mesenchymal stem cells: *Stem Cells Dev* 19: 423-438.
3. Russo E, Caprnda M, Kruzliak P, Conaldi PG, Borlongan CV, et al. (2022) Umbilical Cord Mesenchymal Stromal Cells for Cartilage Regeneration Applications. *Stem Cells Int* 245: 41-68.
4. Serrenho I, Rosado M, Dinis A, M Cardoso C, Graos M, et al. (2021) Stem Cell Therapy for Neonatal Hypoxic-Ischemic Encephalopathy: A Systematic Review of Preclinical Studies. *Int J Mol Sci* 22: 3142.
5. Liu Y, Fang J, Zhang Q, Zhang X, Cao Y, et al. (2020) Wnt10b-overexpressing umbilical cord mesenchymal stem cells promote critical size rat calvarial defect healing by enhanced osteogenesis and VEGF-mediated angiogenesis. *J Orthop Translat* 23: 29-37.
6. Arrigoni C, Arrigo D, Rossella V, Candrian C, Albertini V, et al. (2020) Umbilical Cord MSCs and Their Secretome in the Therapy of Arthritic Diseases: A Research and Industrial Perspective. *Cells* 9: 13-43.
7. Kohler H, Pashov AD, Kieber-Emmons T (2019) Commentary: Immunology's Coming of Age. *Front Immunol* 10: 21-75.
8. Leone P, Solimando AG, Malerba E, Fasano R, Buonavoglia A, et al. (2020) Actors on the Scene: Immune Cells in the Myeloma Niche. *Front Oncol* 10: 597-598.
9. Kageyama T, Yoshimura C, Myasnikova D, Kataoka K, Nittami T, et al. (2018) Spontaneous hair follicle germ (HFG) formation in vitro, enabling the large-scale production of HFGs for regenerative medicine. *Biomaterials* 154: 291-300.
10. Kageyama T, Chun YS, Fukuda J (2021) Hair follicle germs containing vascular endothelial cells for hair regenerative medicine. *Sci Rep* 11: 6-24.