



Combining Machine Learning and Heart Rate Variability Data to Predict 爛

Citation: Sudhindra (2022) Combining Machine Learning and Heart Rate Variability Data to Predict GABA Receptor Antagonist-Induced Convulsion in Cynomolgus Monkeys. *World J Pharmacol Toxicol* 5: 145.

Drug-induced convulsion may be a severe adverse event; but,

Copyright: © 2022 Sudhindra. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

information. M summarizes the number of convulsion alarms issued by the models of M1-M4 once PTX was dosed. The 24-h data contained important changes within the activities of ANS because of feeding and sleeping; HRV data containing such changes weren't applicable for model training [5]. Thus, the convulsion prediction model has got to be trained from HRV data throughout resting conditions, during which ANS activities don't considerably fluctuate. This indicates that the modelling data have to be selected carefully, which is a common problem in machine learning.

References

1. Wong CH, Siah KW, Lo AW (2019) Estimation of clinical trial success rates and related parameters. *Biostatistics* 20: 273-286.
 2. Tsuji H, Larson MG, Venditti FJ, Manders ES, Evans JC, et al. (1996) Impact of reduced heart rate variability on risk for cardiac events. The Framingham heart study. *Circulation* 94: 2850-2855.
 3. Shafer F, Ginsberg JP (2017 of h) An overview eart rate variability metrics and norms. *Front Public Health* 5: 258.
 4. Kim HG, Cheon EJ, Bai DS, Lee YH, Koo BH (2018) Stress and heart rate variability: A meta-analysis and review of the literature. *Psychiatry Investig* 15: 235-245.
 5. Kerem DH, Geva AB (2005) Forecasting epilepsy from the heart rate signal. *Med Biol Eng Comput* 43: 230-239.
-