



## Dual Malignancy of Pancreas: A Rare Case Report and Review of Literature

Aniq Gajdhar<sup>1\*</sup>, Selvakumar Balakrishnan<sup>1</sup>, Anand Vijai<sup>1</sup>, Palanisamy Nalankilli<sup>1</sup>, Basavaraju Srikanth<sup>2</sup> and Chinnusamy Palanivelu<sup>1</sup>

<sup>1</sup>Department of Hepato-Pancreato-Biliary and Minimal Access Surgery, GEM Hospital and Research Center, Coimbatore, Tamil Nadu, India

<sup>2</sup>Department of Radiology, GEM Hospital and Research Center, Coimbatore, Tamil Nadu, India

---

**\*Corresponding author:** Aniq Gajdhar, GEM Hospital and Research Center, 45, Pankaja Mills Road, Ramanathapuram, Tamil Nadu, India – 641045, E-mail: aniq.gajdhar@gmail.com

**Received:** 7-Jan-2022, Manuscript No: JCD-22-51319, **Editor assigned:** 10-Jan-2022, PreQC No: JCD-22-51319(PQ), **Reviewed:** 24-Jan-2022, QC No: JCD-22-51319, **Revised:** 28-Jan-2022, Manuscript No: JCD-22-51319(R), **Published:** 4-Feb-2022, DOI: 10.4172/2476-2253.1000136

**Citation:** Gajdhar A, Balakrishnan -22-

with at least two different malignant components, without mixed or transitional area [2]. They can occur in any organ of the body with stomach and esophagus being the most commonly involved organs. Pancreas is a very unusual site for such presentation.

As per the available literature, sporadic cases comprising of PDA with IPMN, IPMN with NET, solid pseudo papillary neoplasm with NET and cancer of bile duct with pancreas have been reported [3,4]. Chang et al. [5] reported one such case of solitary concomitant NET and PDA in a 58-year-old lady and provided a simple classification for such neoplasms. Similarly, a case of pancreatic collision tumor