problems. Since prothrombin time (PT) and partial thromboplastin time (PTT) are used to assess the extrinsic and intrinsic pathway respectively, these two measures can be helpful in detecting possible problems in the coagulation cascade.

So we hypothesized that Orlistat, by means of inhibiting the absorption of lipids, can potentially reduce the absorption of vitamin K and possibly making some dysfunctions in the coagulation cascade. Therefore, we decided to give Orlistat to Wistar rats and measure their PT and PTT for assessing any possible coagulation dysfunction.

## Materials and Methods

54 male Wistar rats, with the mean age of 8 weeks and mean weight of 300 grams were obtained from the animal house of the Shahid

| | | | |
|---|---|---|---|
| Control 1 vs. Control 2 (One Month) | 0.5 | -28.59 to 29.59 | No |
| Control 1 vs. Case (One Month) | -5.333 | -54.09 to 43.42 | No |
| Control 2 vs. Case (One Month) | -5.833 | -38.39 to 26.72 | No |
| Control 1 vs. Control 2 (Three Months) | 4.167 | -18.23 to 26.57 | No |
| Control 1 vs. Case (Three Months) | 7.833 | -4.976 to 20.64 | No |
| Control 2 vs. Case (Three Months) | 3.667 | -14.70 to 22.03 | No |
| Case (One Week) vs. Case (One Month) | -9.667 | -51.40 to 32.07 | No |
| Case (One Week) vs. Case (Three Months) | 2.5 | -9.902 to 14.90 | No |
| Case (One Month) vs. Case (Three Months) | 12.17 | -33.55 to 57.88 | No |

**Table 2:** Results of Tukey's test for PTT.

Also a comparison was made between means of PT and PTT in different time periods which can be seen in Figures 1 and 2, respectively.

**Figure 1:** PT Change Pattern with time.

## Discussion and Conclusion

Our research shows that Orlistat increases PT in every three experiments, whereas alcohol decreased PT in the one week experiment and increased PT in the one month and three months experiments. The pattern of PT change with alcohol consumption didn't follow a predictable pattern through time, thfollow time, t' '

So because of the reasons mentioned above, we decided to test our hypothesis on Wistar rats. These rats have less genetic variability among their population, so the size of the sample doesn't need to be large. Also, the results of this experiment are fairly distributable to the human population since 90% of the rat genome is similar to humans [11].

During the time period of our study (3 months), we couldn't find any significant increase in PT or PTT due to Orlistat consumption. However, the significancy of PT and PTT change, increased more as the time of Orlistat consumption got longer. These results make the probability those longer times.

(>3 months) of Orlistat usage may lead to significant increase in PT and PTT.

The mentioned evidences suggest that Orlistat as an FDA approved anti-obesity drug doesn't cause any serious coagulopathies (if any coagulopathy happens), although theoretically it can cause vitamin K malabsorption and thus causing increases in both PT and PTT. But it should be taken to account that people who use other vitamin K inhibitors such as Warfarin, should use Orlistat with care.

## Ethics

All the steps of this research were in accordance to European convention for the protection of vertebrate animals used for