

Effect of Vestibular Stimulation Versus Whole Body Vibration on Standing Balance in Children with Spastic Diplegic Cerebral Palsy

Ashish Parashar*, Monalisa Pattnaik and Patitapaban Mohanty

Swami Vivekanand National Institute of Rehabilitation Training and Research, Olatpur, Bairoi, Cuttack, India

*Corresponding author: Ashish Parashar, Swami Vivekanand National Institute of Rehabilitation Training and Research, Olatpur, Bairoi, Cuttack, India, Tel: 919753202681; E-mail: ashish.parashar2681@gmail.com

Received date: March 17, 2017; Accepted date: April 24, 2017; Published date: May 01, 2017

Copyright: © 2017 Parashar A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Introduction: Most cerebral palsy children have deficits in balance, co-ordination, and gait throughout childhood and adulthood. So, it is essential to seek an ideal physical therapy program to help in solving such widespread problem. The present study was conducted to compare between the effect of whole body vibration and vestibular stimulation on standing balance in children with spastic diplegic cerebral palsy.

Method: 30 children were randomly assigned to group 1 (n=15, male: 11, female: 4, mean age: 6.2, SD 2.48) and group 2 (n=15, male: 8, female: 7, mean age: 6.9, SD 2.01). Balance was measured by Pediatric balance scale and parents perception on balance. Group 1 received vestibular stimulation in addition to conventional physiotherapy while group 2 received whole body vibration addition to the same physical therapy program given to the group 1.

Statistical analysis: Data was analyzed using Mann Whitney U test by taking the change of score from pre to post. To find out the difference between the groups for changed score of Pediatric balance scale. Independent T-test was used for Parents perception on balance and within group difference analysis was done using Wilcoxon Signed Rank test for both the group separately.

Results: The Group 1 and Group 2 showed an change of 21.10 and 9.90 ranks on PBS and 3.73 & 1.60 on

nerve Vfg and activates monosynaptic and polysynaptic fY Y arcs resulting in contraction of muscle

Subconscious contraction of the muscles causes many more muscle Vfg to contract than in a conscious, voluntary movement [19] fY Y through the heightened EMG activity [20,21].

Y Y of sustained vibratory stimulation on muscle contraction, posture and kinesthetic perceptions are much more complex than merely contraction of the muscle being vibrated.

Vestibular system is one of the main structures to maintain balance, as it serves as an absolute reference in relation to the others, such as visual and somatosensorial systems. Adequately administered vestibular stimulation has been reported to improve balance; the vestibular-spinal fY Y generates body motion compensation, to maintain head and postural stability, and thus preventing falls [22].

Vestibular system when stimulated elicits responses that facilitate, muscles, to evoke movements of head, trunk and limbs to compensate for postural sway. Y cerebellum with input from the vestibular system provides balance, orientation & coordination [23].

Vestibular stimulation over a period of time would improve sensory integration and balance in individual with cerebral palsy [24]. Vestibular stimulation could enhance arousal level, visual exploratory behavior; motor development and balance and fY Y integration in infants who are at risk and in development delay disorders [25].

Research has concluded the Y Y of SI therapy & particularly vestibular stimulation for balance was Y Y in children with CP [26].

A comparative study between Suspension therapy and whole body vibration was done for improving standing balance in children with cerebral palsy. G[b] Wbh X YfYbW was observed in favor of suspension therapy which was attributed to the Y Y of spider cage on improving function of vestibular system through development of equilibrium reactions to maintain & regain balance during standing.

Application of vestibular stimulation for CP children led to g[b] Wbh improvement in standing balance. Also, the children showed a new strategy which would lead to an increase in exploratory behavior which would help them to maintain balance.

Aim of study. Y aim of the study was to compare the Y Y of vestibular stimulation with whole body vibration on balance in children with spastic diplegic cerebral palsy.

Methodology

Study design

A pre-test post-test experimental study design

Sample size: 30 children randomly allocated to 2 Groups by chit picking

Group 1: Conventional therapy and vestibular stimulation (n=15, male: 11, female: 4, mean age: 6.2, SD 2.48)

Group 2: Conventional therapy and Whole body vibration (n=15, male: 8, female: 7, mean age: 6.9, SD 2.01)

Study duration: 4 weeks

Inclusion criteria

Children diagnosed as spastic diplegic cerebral palsy, of either sex in the age group ranging from 3 to 12 years, could understand and follow commands given by the therapist and could stand independently or with one hand support for at least 10 seconds.

Exclusion criteria

Children with types of cerebral palsy other than spastic diplegia, History of convulsions/epilepsy, impaired cognitive function, negative tilt board tip test, any recent surgical procedure for correction of deformity; g: tissue release etc.,

Procedure

Informed consent was obtained from parents of children who Z` ``X inclusion and exclusion criteria and randomly assigned to Group 1 and Group 2. Pre-test measurements of the dependent variables pediatric balance scale (PBS) and parents perception on balance (PPB) were measured on recruitment for the study by one investigator, who was blind to the type of treatment given.

Paediatric Balance Scale (PBS) is a reliable measure of functional balance for school age children with mild to moderate motor impairments. Test retest reliability=0.87 to 1.0, interrater reliability=0.997, correlation=0.89 to 1.0 [55].

Parent's perception on balance (PPB) is an observational measure of functional balance of child by parents using Visual Analogue Scale. Parents were asked to grade child's standing balance on 0-10 scale.

Group 1 treated by conventional physiotherapy + Vestibular stimulation

Technique of application of Vestibular stimulation: Procedure was explained to the parents. SPINNING- Child was asked to sit on hammock and spun 10 times to the right and 10 times to the Y Z SLIDING- Child was asked to lie in supine position on a large physio roll, then the child was accelerated forward and backward direction 10 times by holding both the legs. BOUNCING UP AND DOWN- Child was made to sit on a large physio roll, holding the child from pelvis by therapist then bounced the child vertically up and down for 5 minutes. Total treatment duration was 10 minutes.

Group 2 treated by conventional physiotherapy + whole body vibration

Technique of application of Whole body vibration

Both the groups received conventional physiotherapy which includes bridging, quadruped, kneeling, sit to stand, reaching. Treatment was given for 5 days a week for 4 weeks.

Data Analysis

Data was analysed using Mann Whitney U test by taking the change of score from pre to post. To compare the X² test between the groups for changed score of Paediatric balance scale. Independent T-test was used for Parents perception on balance and within group X² test analysis was done using Wilcoxon Signed Rank test for both the group separately (Figure 1).

An alpha level of 0.05 of significance was set for both the tests.

Analysis was performed using SPSS package 23 version.

Results

Study demonstrated Y YW

33. Hosseini SA, Haghighi HA, Zadeh BZ, Pishyareh E, Talebian S (2012) Vestibular stimulation on quiet stance balance in 3- to 7-year-old children with cerebral palsy.
34. Damiano DL, Abel MF (1998) Functional outcomes of strength training in spastic cerebral palsy. *Arch Phys Med Rehabil* 79: 119-125
35. Wiley ME, Damiano DL (1998) Lower-extremity strength deficit in spastic cerebral palsy. *Dev Med Child Neurol* 40: 100-107.
36. Roelants M, Delecluse C, Verschueren S, Levin O, Stijnen V (2006) Whole body vibration induced increase in leg muscle activity during X squat exercises. *Journal of strength and conditioning research* 20: 124-129
37. Saggini R, Vecchiet J, Iezzi S, Racciatti D, Uhl G, et al. (2006) Submaximal aerobic exercise with mechanical vibrations improves the functional status of patients with chronic fatigue syndrome. *Eura Medicophys* 42: 97-102
38. Fagnani F, Giombini A, Di Cesare A, Pigozzi F, Di Salvo V (2006) Effect of a whole-body vibration program on muscle performance and strength in female athletes. *Am J Phys Med Rehabil* 85: 956-962
39. Mahieu NN, Witvrouw E, Van de VD, Michilsens D, Arbyn V, et al. (2006) Improving strength and postural control in young skiers: whole-body vibration versus equivalent resistance training. *J Athl Train* 41: 286-293
40. Roelants M, Delecluse C, Verschueren SM (2004) Whole-body-vibration training increases knee-extension strength and speed of movement in older women. *Journal of the American Geriatrics Society* 52: 901-908
41. Roelants M, Delecluse C, Goris M, Verschueren S (2004) Effect of 24 weeks of whole body vibration training on body composition and muscle strength in untrained females. *Int J Sports Med* 25: 1-5
42. Verschueren SM, Roelants M, Delecluse C, Swinnen S, Vanderschueren D, et al. (2004) Effect of 6-month whole body vibration training on hip density, muscle strength, and postural control in postmenopausal women: a randomized controlled pilot study. *Journal of bone and mineral research* 19: 352-359
43. Torvinen S, Kannus P, Sievänen H, Järvinen TA, Pasanen M, et al. (2002) Effect of a vibration exposure on muscular performance and body balance. Randomized cross-over study. *Clin Physiol Funct Imaging* 22: 145-152
44. Torvinen S, Kannus P, Sievänen H, Järvinen TA, Pasanen M, et al. (2002) Effect of four-month vertical whole body vibration on performance and balance. *Med Sci Sports Exerc* 34: 1523-1528
45. Wilcock IM, Whatman C, Harris N, Keogh JW (2009) Vibration training: could it enhance the strength, power, or speed of athletes? *J Strength Cond Res* 23: 593-603
46. Luo J, McNamara B, Moran K (2005) Use of vibration training to enhance muscle strength and power. *Sports Med* 35: 23-41.
47. Ebersbach G, Edler D, Uebachs O, Wissel J (2008) Whole body vibration versus conventional physiotherapy to improve balance and gait in Parkinson's disease. *Arch Phys Med Rehabil* 89: 399-403
48. Kawanabe K, Kawashima A, Sashimoto I, Takeda T, Sato Y, et al. (2007) Effect of whole-body vibration exercise and muscle strengthening balance, and walking exercises on walking ability in the elderly. *Keio journal of medicine* 56: 28-33
49. Bogaerts A, Verschueren S, Delecluse C, Claessens AL, Boonen S (2007) Effect of whole body vibration training on postural control in older individuals: a 1 year randomized controlled trial. *Gait Posture* 26: 309-316
50. Bautmans I, VanHees E, Lemper JC, Mets T (2006) Feasibility of Whole Body Vibration in institutionalised elderly persons and its effect on balance.