



# & WBMVBUJPO PG UIF & GGJD #WQDBZ NP QP SBT D I 1 IZUP SFNFEJBUJPO "HFOU JO 8BTUF XBUFS 5S

Irfana Showqj\*, Farooq Ahmad Lone and Javeed Iqbal Ahmad Bhat

Division of Environmental Sciences, Sheri-Kashmir University of Agricultural Sciences and Technology of Kashmir, Shalimar, Jammu and Kashmir, India

## Abstract

,Q WKH SUHVHQW VWXG\ WKH HI¿FLHQF\ RI GXFNZHHG Lemna minor (L.) as an effective na  
wastewater treatment was examined in an outdoor aquatic system. Duckweed plants were inoculated into wastewater  
DQG WDS ZDWHU V\WHPV IRU WUHDWPHQW RYHU ¿IWHHQ GD\¶V UHWHQWLRQ SHU  
VDP SOHV ZHUH WDNHQ EHORZ GXFNZHHG FRYHU DIWHU ¿IWHHQ GD\ WR DVVH  
water from different pollutants. For comparison, the plants were also grown in tubs containing tap water. The results  
show that concentrations of Nitrogen (N), Phosphorus (P), Potassium (K), Calcium (Ca), Magnesium (Mg), Copper  
(Cu), Zinc (Zn), Nickel (Ni), Chromium (Cr), Cadmium (Cd) and Lead (Pb) decreased by 93.4%, 99.9%, 93.9%,  
98.5%, 91.9%, 85.0%, 95.0%, 90.0%, 99.8%, 99.5% and 95.0% respectively in waste water and subsequently  
these elements exhibited an increasing concentration in the plant body. Almost similar results were obtained when  
the plants were grown in tubs containing tap water. Biochemical parameters viz. chl-a, b, total-chl, carbohydrates  
and proteins as well as nutrient status of the macrophyte increased after the completion of the retention period both  
LQ ZDVWH DQG WDS ZDWHU 5HVXOWV FRQ¿UP WKDW GXFNZHHGV FDQ HIIHFWLYH

**Keywords:** Lemna minor, Wastewater; Retention period

## Introduction

Multiple environmental factors in association with anthropogenic activities have significantly altered our aquatic ecosystems. Over use of chemical fertilizers and intensification of industrial activities

for the removal of contaminants from the environment are cheaper than the conventional remediation technologies [1]. In recent years, there has been a growing struggle to provide efficient, inexpensive, and environmentally friendly options for the remediation of trace elements and other contaminants in waste water. Moreover, aquatic plants are of special interest unlike the terrestrial plants, because they are capable of bioconcentrating many nutrients and heavy metals in large quantities [2-4]. In Kashmir valley Anchar lake is one of the lakes

\*Corresponding author: Irfana Showqj, Division of Environmental Sciences, Sheri-Kashmir University of Agricultural Sciences and Technology of Kashmir, Shalimar, Jammu and Kashmir, India, Tel: +919796598837; E-mail: irfanashowqj@gmail.com

Received June 29, 2017; Accepted July 24, 2017; Published July 26, 2017

Citation: Showqj I, Lone FA, Bhat JIA (YDOXDWLRQ RI WKH Duckweed (Lemna minor L.) as a Phytoremediation Agent in Wastewater Treatment in Kashmir Himalayas. J Bioremediat Biodegrad 8: 405. doi: 10.4172/2155-6199.1000405

Copyright: © 2017 Showqj I, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

reduction percentage of 85.0%, 95.0% and 90.0% respectively (Table 1). On the other side, these micronutrient (Cu, Zn and Ni) show high concentration in the plant body during the retention time with an increasing percentage of 34.69%, 69.44%, and 66.67% respectively in the plant body (Table 2). Similar results have been observed in tap water that was used for comparison purposes where micronutrients zinc and copper had decreased in concentration after *Lemna minor* grown in it for 15 days with a reduction percentage 100% and subsequently in plant body these nutrients have shown increasing percentage of 2.22% and 0.71% respectively (Table 2) however nickel has shown no change as it was not observed in tap water during initial analysis. In the line of present study Wa a et al. [17] reported that duckweed treatment system reduced zinc by 93.6% and copper by 100% during eight days retention period. Zaltauskaite et al. [15] also found that Zn was most efficiently removed, depending on the initial Zn concentration *Lemna minor* removed between 42.3-77.8% of Zn. They also found that after the 7 days of elements exposure to *Lemna minor* treatment, the concentration nickel was reduced significantly.

The major cause of micronutrient reduction in waste water / tap water was utilization by the *Lemna minor* plant for body formation and development. Duckweeds require a number of macro and micronutrients for their normal growth. Nutrients are absorbed through all surfaces of the duckweed leaf. DWRP [18] reported that the highest growth rate for Lemnaceae under optimal laboratory conditions is about 0.66 generations per day that is equal to a doubling time of 16 hours. It has been reported that duckweeds generally double their mass in 16 hours to 2 days under optimal conditions which further helps in uptake of nutrients from the growing media for developing body tissue. As also reported by Korner and Vermaat [19] that the nutrients removed by duckweeds from growing media are mainly realized by newly grown tissue of the plants. Duckweed has a high mineral absorption capacity and can tolerate high organic loading as well as high concentrations of micronutrients.

#### Heavy metals (Cd, Cr and Pb)

The concentrations of heavy metals such as Cd, Cr and Pb reduced after *Lemna minor* was grown on wastewater for 15 days retention period. These metals have reached their minimum concentration of 0.001, 0.001 and 0.01 mg/l respectively with a reduction percentage of 99.5%, 99.8% and 95.0% respectively, however in plant body their concentration has increased during the study period with an increasing percentage of 92.06% 43.59% and 0.52% respectively. In case of tap water cadmium and lead were not observed during initial analysis however chromium had decreased in concentration from 0.20 ppm to 0.00 ppm 73 (, h)4 (o)16 (w)8 (e)-8 (v)84 Tw T\* [(pcs)-8 (er)-29 (v)tio 327.4

in plant body chromium have shown increasing percentage of 3.23%, whereas cadmium and lead have shown no change in their initial concentrations (Table 2). In concurrence with the present findings, Wa a [17] studied that duckweed aquatic treatment system performed 100% lead and 66.7% of cadmium removal from wastewater after 8 days treatment period. Similar results have also been found by Kara et al. [20] and concluded that duckweed *Lemna minor* take up Pb, Cu, Fe, Cd and Ni from contaminated solutions. Removal of heavy metals from

2. Keskinan O, Goksu MZL, Basibuyuk M, Forster CF (2004) Heavy metal adsorption properties of a submerged aquatic plant (*Ceratophyllum demersum*). *Bioresource Technology* 92: 197-200.
3. Vardanyan LG, Ingole B (2006) Studies on heavy metal accumulation in aquatic macrophytes from Sevan (Armenia) and Carambolim (India) systems. *Environmental International* 32: 208-218.
4. Pratas J, Favas PJC, Paulo C, Rodrigues N, Prasad MNV (2012) Uranium accumulation by aquatic plants from uranium-contaminated water in Central Portugal. *International Journal of Phytoremediation* 14: 221-234.
5. 3DU 3HUNLQV 5\* 0DVRQ &) 5HGXFWRQ LQ SKRWRV\QWKHWLF HI FLHQF\ of *Cladophora glomerata*, induced by overlying canopies of *Lemna* spp. *Water Research* 36: 1735-1742.
6. American Public Health Association (1992) *Standard Methods for the Examination of Water and Wastewater*. Washington, USA.
7. Strain HH, Cope BT, Walter AS (1971) Analytical procedures for the isolation, LGHWL FDLRQ HVWLPDLRQ DQG LQYHVWLDWRQ RI WKH FKORURSK\OOV 0HWKRGV LQ *Enzymology*, Academic Press, New York, USA 23: 452-457.
8. Duxbury AC, Yentsch CS (1956) Plankton pigment monographs. *Journal of Marine Research* 15: 91-101.
9. Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F (1956) Colorimetric