



**Keywords:** Decision recommendation; Feature interpretation;  
Stacked ensemble

## Introduction

ere was once an enormous main-e ect of age for all hallucination



- 
2. Waters F, Woods A, Fernyhough C (2014) Report on the 2nd International Consortium on Hallucination Research: evolving directions and top-10 "hot spots" in hallucination research. *Schizophr Bull* 40:24-27.
  3. Sartorius N, Jablensky A, Korten A, Embregt G, Anker M, et al. (1986) Early manifestations and first-contact incidence of schizophrenia in different cultures. A preliminary report on the initial evaluation phase of the WHO Collaborative Study on determinants of outcome of severe mental disorders *Psychol Med* 16:909-28.
  4. Harding AJ, Broe GA, Halliday GM (2002) Visual hallucinations in Lewy body disease relate to Lewy bodies in the temporal lobe 125:391-403.
  5. Sarah HL, Taylor J, Sarah K, Emma E, Jill MH, et al. (2021) Psychosis proneness, loneliness, and hallucinations in nonclinical individuals 16: 0251753.
  6. Flavie W, Charles F (2017) Hallucinations: A Systematic Review of Points of Similarity and Difference Across Diagnostic Classes. *Schizophr Bull* 43: 32-43.
  7. Eguchi A, Nomura S, Gilmour S, Harada N, Sakamoto H, et al. (2021) Suicide by gender and 10-year age groups during the COVID-19 pandemic vs previous five years in Japan: an analysis of national vital statistics. *Psychiatry Research* p.114173.
  8. Isumi A, Doi S, Yamaoka Y, Takahashi K, Fujiwara T, et al. (2020) Do suicide rates in children and adolescents change during school closure in Japan? The acute effect of the first wave of COVID-19 pandemic on child and adolescent mental health. *Child Abuse & Neglect* 110:104680.
  9. Nomura S, Kawashima T, Yoneoka D, Tanoue Y, Eguchi A, et al. (2021) Trends in suicide in Japan by gender during the COVID-19 pandemic, up to September 2020. *Psychiatry Research* 295:113622.
  10. Pirkis J, John A, Shin S, Arya V, Analuisa AP, et al. (2021) Suicide trends in the early months of the COVID-19 pandemic: an interrupted time-series analysis of preliminary data from 21 countries. *The Lancet Psychiatry* 7:579-588.