


**Diagnosis**



usual diagnosis of genital herpes is based on a clinical history of genital ulcers and a positive culture of the ulcers. The diagnosis is usually confirmed by a positive culture of the ulcers. The diagnosis is usually confirmed by a positive culture of the ulcers.

- 
1. Tarimo CS (2019) Prevalence and predictors of failure in labor induction among pregnant women delivered in Northern-Tanzania 2000-2015: A Registry-based Retrospective Cohort Study. *Tanzan Med J* 30: 13-36.
  2. Lyndrup J, Legarth J, Weber T, Nickelsen C, Guldbæk E (1992) Predictive value of pelvic scores for induction of labor by local PGE<sub>2</sub>. *Eur J Obstet Gynecol Reprod Biol* 47: 17-23.
  3. Khan NB, Ahmed I, Malik A, Sheikh L (2012) Factors associated with failed induction of labour in a secondary care hospital. *J Pak Med Assoc* 62: 6.
  4. Tripathy P, Pati T, Baby P, Mohapatra SK (2016) Prevalence and predictors of failed. *Int J Pharm Sci Rev Res* 39: 189-94.
  5. Mbukani R, Kakoma J (2012) Is Nulliparity A Risk Factor For Poor Obstetrical And Neonatal Outcomes In Rwandan District Hospitals? A Prospective Observational Study at Muhima District Hospital. *Rwanda Med J* 69: 50-53.
  6. ) Impact of labor induction, gestational age, and maternal age on cesarean delivery rates. *Obstet Gynecol* 102: 287-293.
  7. Verhoeven CJ, Van Uytrecht CT, Porath MM, Mol BWJ (2013) Risk factors for caesarean delivery following labor induction in multiparous women. *J Pregnancy* 2013: 820892.
  8. Hannah ME (1993) Post term pregnancy: should all women have labour induced ? A review of the literature. *Fetal Matern Med Rev* 5: 3-17.
  9. Galal M, Symonds I, Murray H, Petraglia F, Smith R (2012) Post term pregnancy. *Facts, Views & Vis Obstet Gyn* 4: 175.
  10. Obstetricians ACo, Gynecologists (2009) ACOG practice bulletin no. 107: Induction of labor. *Obstet Gynecol* 114: 386-397.
-