



***Corresponding author:** Muluneh Mekiso Halengo, Worabe Agricultural Research Centre, Ethiopia, Tel: +251 986 945 389; Email: mekisomh@gmail.com

Received date: August 17, 2020; **Accepted date:** August 21, 2020; **Published date:** August 29, 2020

Citation:

Citation: : Muluneh Mekiso Halengo(2020); Genotype by Environment interaction of food barley (*Hordeum vulgare* L.) genotypes in Southern Ethiopia; J Plant Genet Breed S (1): 110

Location	Altitude (masl)	Mean annual rainfall (mm)*	Average temperature (°C)*	Soil texture	Global position	
					Latitude	Longitude
Mierab-						

Citation: : Muluneh Mekiso Halengo(2020); Genotype by Environment interaction of food barley (*Hordeum vulgare* L.) genotypes in Southern Ethiopia; J Plant Genet Breed S (1): 110

Genotype name	Mean grain yield (t ha ⁻¹)	Wi	Pi	CV	(bi)	(S ² d)	ASV
G1(HB-1307)	3.79	4.45	0.37				


AMMI1 Biplot Analysis

the variety had the largest positive interaction with that specific

In GGE biplot for grain yield of food barley varieties, the first and captured 92.2% of the G+GE sum of square, it means that different

described the significant GEI in wheat by AMMI biplot analysis.

these varieties had specific adaptations.



Citation: : Muluneh Mekiso Halengo(2020); Genotype by Environment interaction of food barley (*Hordeum vulgare* L.) genotypes in Southern Ethiopia; J Plant Genet Breed S (1): 110

References:

- a. Symposium, Alexandria, Egypt. ICARDA, Aleppo, Syria. pp xii +