”º •Ãº½Á¶· ³Â ¯º •˜¯À Â¶ ¦±· –

Open Acces

GIS-based Framework for the Simulation of the Impacts of Sea Level Rise and Coastal Flooding on Oman

Al-Buloshi A, Al-Hatrushi S* and Charabi Y

Department of Geography, Sultan Qaboos University, Muscat, Sultanate of Oman

Keywords Oman; SLR scenarios; GIS; Coastal vulnerability

Introduction

By 2100, the 2001 IPCC ird Assessment Report (TAR) [1] predicted a rise in Sea level between 20 and 70 cm. However, in the 2007 IPCC [2] AR4 projected that sea level will rise from 18-59 cm during the same period. A recent study published by Rahmstorf, Foster, and Cazenave [3] (RFC 12) consider the above IPCC projection as too conservatives and underestimate future SLR, due to the limitations of the IPCC model, because they did not involve dynamic ice melting processes.

RFC12 uses a semi-empirical approach based on the proportionality between global warming and SLR. Further warming will lead to more ice melt and reinforcing the melting cycle. RFC12 concludes that satellite-based linear trend 1993–2011 is 3.2 ± 0.5 mm yr^{-1} which is 60% faster than the best IPCC estimate of 2.0 mm yr^{-1} for the same interval.

Most likely, the di erence between RFC12 semi-empirical

*Corresponding author: Al-Hatrushi S, Department of Geography, Sultan Qaboos University, Muscat, Sultanate of Oman, Tel: +968-24142036; E-mail: hatrushi@squ.edu.om

Received Jun 30, 2014; Accepted November 11, 2014; Published November 20, 2014

Citation: Al-Buloshi A, Al-Hatrushi S, Charabi Y (2014) GIS-based Framework for the Simulation of the Impacts of Sea Level Rise and Coastal Flooding on Oman. J Earth Sci Clim Change 5: 238. doi:10.4172/2157-7617.1000238

Copyright: © 2014 Al-Buloshi A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

shoreline has long been recognized as a problem of growing magnitude. At present, coastal erosion and accretion occurs as natural processes



Figure 2: Topography of Oman.



Figure 3: Built-up population areas in Oman, 2007.

e Tidal Datum and Reference Benchmarks data based was obtained from the Oman National Hydrographic O ce. e assumed tidal range in the Sea of Oman was 0.89 to 1.70 meters. Since 1990, Chart Datum has been established at over 40 sites along Oman's coast from observations conducted by, or under the supervision of, the Royal Navy of Oman's Hydrographic Department [9-12]. A total of 7 SLR scenarios were considered: 0.2 meters, 0.5 meters, 1 meter, 2 meters, 3 meters, 4 meters and 5 meters.

To assess the maximum inundation risk to coastal communities, these levels were added to Mean High Tide (MHT) levels. e sea level scenarios were overlain on land use/elevation datasets to estimate the inundation risk among all land use categories. e coastal line of the governorate of Dhofar was excluded from the estimation of the risk due to the lack of appropriate land use datasets. All the grid cells that would be inundated based on a user-de ned horizontal spatial resolution of 0.5 meters for the entire coastline.

Results and Discussions

e results of the assessment show that Oman is highly vulnerable