

Headache Attributed to Airplane Travel in a Child-First Case in Brazil

Luara Lis Barbosa Boson¹, Lucas Aguiar Alencar Oliveira², Fernando Jacó Silva Moreira³, Mariana Leite Pereira³, Rayssa Fernandes Souza Coelho³, Allyson Coelho Ribeiro⁴, Augusto César Beserra Martins⁴, José Hugo Andrade Santos Dantas⁴, Isadora Costa Coelho Gayoso-Almendra⁴, Ítalo Araújo Rios Brandão⁴, July Lima Gomes⁴, Kamilla Gomes Sales Souza⁴, Laysa Moura Cardoso Leal⁴, Luís Gustavo Silva Bacelar Andrade⁴ (g ðÖ8Å I Av L S•Â H L` d& ð°

T lg

A.	At least two episodes of headache fulfilling criterion C
B.	The patient is travelling by aeroplane
C.	Evidence of causation demonstrated by at least two of the following:
	1. Headache has developed exclusively during aeroplane travel
	2. Either or both of the following:
	a) headache has worsened in temporal relation to ascent after take-off and/or descent prior to landing of the aeroplane
	b) headache has spontaneously improved within 30 minutes after the ascent or descent of the aeroplane is completed
	3. Headache is severe, with at least two of the following three characteristics:
	a) unilateral location
	b) orbitofrontal location (parietal spread may occur)
	c) jabbing or stabbing quality (pulsation may also occur)
D.	Not better accounted for by another ICHD-3 diagnosis.

fgh

T lg

f \

]b aa a aljcb

