

Helicobacter Pylori Infection and Gallbladder Diseases: a Mini-review


Mohammad Javad Yavari Barhaghtalaba*1 Mohammad Bagher Jahantaba,2 Amir Abbas Safaripoura,2 Reza Hosseinpoura,2 Saadat Mehrabi Sisakhta, Cambyz Irajieb2

FÖ^}æ/c { ^}ç/[-hÖ^}^æ|hÜ~i*^i^ÉÅYæ•~hW)âç^i•æ~h[-hT^â&æ|hÜ&â^}&^ÉÅYæ•~iÉh/iæ}

GÖ^}æ/c { ^}ç/[-hT^â&æ|hÜ&â^} [i [*ÉhÜ@â/iæ:hW)âç^i•æ~h[-hT^â&æ|hÜ&â^}&^ÉÅYæ•~iÉh/iæ}

*Corresponding author: Mohammad Javad Yavari Barhaghtalaba, Department of General Surgery, Yasuj University of Medical Sciences, Yasuj, Iran, Email- mj.yavari.barhaghtalab@gmail.com

BahM:ipaloo m



Gallbladder cancer

Risk factors for the development of gallbladder cancer are patient demographics, gallbladder abnormality, patient exposure, and Salmonella and Helicobacter infections. The association between H[21 22]. pylori infection and biliary tract carcinoma (gallbladder cancer,