

Keywords: ...

Introduction

...

...

Discussion

...

Pari Mane, Department of Radiology, University of Zhejiang, China, E-mail: Parimane54@gmail.com

01-Apr-2023, Manuscript No. jpch-23-91008; 07-Apr-2023, PreQC No. jpch-23-91008 (PQ); 21-Apr-2023, QC No. jpch-23-91008; 24-Apr-2023, Manuscript No. Jpch-23-91008(R); 28-Apr-2023, DOI: 10.4172/2376-127X.1000588

... % ...

... % ...

... % ...

Conclusion

... % ...

... % ...

Hydatid cyst of the ovary is a rare type of cystic ovarian lesion. It is caused by the larval stage of the tapeworm *Echinococcus granulosus*. The cyst is usually found in the ovary and is characterized by a thick, fibrous wall and a central core of hyaline fluid. The cyst is usually asymptomatic and is often discovered incidentally during a pelvic examination or imaging study. The diagnosis is usually confirmed by a positive result on a serological test for *Echinococcus* antibodies. Treatment is usually surgical removal of the cyst. The prognosis is generally good, with a low risk of recurrence. However, there is a risk of complications such as rupture of the cyst, which can lead to a severe allergic reaction. Therefore, it is important to be aware of the possibility of this rare condition and to seek medical attention if you experience any symptoms.

Conflict of Interest

1.

Acknowledgement

1.