

Short Communication

Open Access

) Z Q F S B D V T J T J O E V D F E 1 B J O 6 O E F S T U B O E J O H
5 F O T P S 5 Z N Q B O J 4 Z O E S P N F 5 5 5 4 4 Z N Q U P N T

Myriam Westcott *

Dineen Westcott Moore Audiology, Melbourne, Australia

Most clinicians providing tinnitus and hyperacusis therapy agree that hyperacusis can be defined as an abnormal intolerance, heightened sense of volume and physical discomfort in response to certain sounds, which other people can tolerate comfortably. Sounds that are typically difficult to tolerate are loud/impact/sustained sounds, particularly if they are unexpected or in close proximity.

Hyperacusis can be a 8(s)-8(20044)g20044role cardraais3(.)4(d h)22.9(yp)/loTw T* [(33-1.2 Td [(t)-6(h)3(a)(t)-6(4(.5()94)g206 39 cur2(um6)]T2.17(t(erac)-6.9(u)2.9(si)3.1(-3(a)9(n (n a)8(p)-8.9(o)12(t t)6(o t)6(oo)7(l)1d s)-uimi)12.)6(og(u)-5(l)(s p)1t)6(o)7(lerj)9(n)

*Corresponding author: Myriam Westcott, Dineen Westcott Moore Audiology, Melbourne, Australia, Tel: 61 3 94576588; E-mail: myriam@dineenwestcottmoore.com.au

Received December 15, 2015; Accepted February 05, 2016; Published February 09, 2016

Citation: Westcott M (2016) Hyperacusis-induced Pain: Understanding and Management of Tonic Tensor Tympani Syndrome (TTTS) Symptoms. J Pain Relief 5: 234. doi:[10.4172/2167-0846.1000234](https://doi.org/10.4172/2167-0846.1000234)

Copyright: © 2016 Westcott M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

facial muscles in and around the ear and guidance in locating muscle Tensor veli palatini and tensor tympani muscles: anatomical, functional and trigger points in the neck, shoulder and arm. Patients carrying out symptomatic links. *Acta Otorrinolaringol Esp* 61: 26-33.

bs muscle

10. Westcott M (2008) Tonic Tensor Tympani Syndrome -an explanation for everyday sounds causing pain in tinnitus and hyperacusis clients. Poster, IXth International Tinnitus Seminars Gothenburg.
11. Ellenstein A, Yusuf N, Hallett M (2013) Middle ear myoclonus: two informative cases and a systematic discussion of myogenic tinnitus. *Tremor Other Hyperkinet Mov (NY)* 3.
12. Pollak N, Azadarmaki R, Ahmad S (2014) Feasibility of endoscopic treatment of middle ear myoclonus: a cadaveric study. *ISRN Otolaryngol* 2014: 175-268.
13. Westcott M, Sanchez TG, Diges I, Saba C, Dineen R, et al. (2013) Tonic Tensor Tympani Syndrome (TTTS) in Tinnitus and Hyperacusis Patients: A Multi-Clinic Prevalence Study. *Noise Health* 15: 117-128.
14. Riga M, Xenellis J, Peraki E, Ferekidou E, Korres S (2010) Aural symptoms in patients with temporomandibular joint disorders: multiple frequency tympanometry provides objective evidence of changes in middle ear impedance. *Otol Neurotol* 31: 1359-1364.
15. Franz P, Hamzavi JS, Schneider B, Ehrenberger K (2003) Do middle ear muscles trigger attacks of Menière's disease? *Acta Otolaryngol* 123: 133-137.
16. Bance M, Makki FM, Garland P, Alian WA, van Wijhe RG, et al. (2013) Effects of tensor tympani muscle contraction on the middle ear and markers of a contracted muscle. *Laryngoscope* 123: 1021-1027.
17. Curtis AW (1980) Myofascial pain-dysfunction syndrome: the role of nonmasticatory muscles in 91 patients. *Otolaryngol Head Neck Surg* 88: 361-367.
18. Ramírez LM, Ballesteros LE, Sandoval GP (2007) Tensor tympani muscle: strange chewing muscle. *Med Oral Patol Oral Cir Bucal* 12: E96-100.
19. Douglas CR, Avoglio JL, de Oliveira H (2010) Stomatognathic adaptive motor syndrome is the correct diagnosis for temporomandibular disorders. *Med Hypotheses* 74: 710-718.
20. Westcott M (2010) Hyperacusis a clinical perspective on management. *Tinnitus 'L V F R Y H U \ \$ V L D D Q G 3 D F L \ F 7 L Q Q L W X V 6 \ P S R V L X*