alpha1 6.9–8.9 Hz; alpha2 8.9–10.9 Hz; alpha3 10.9–12.9 Hz. Finally, the relative power spectra was computed for each subject in the individual-determined frequency bands. The ratio between the absolute power and the mean power spectra from 2 to 45 Hz engenders the relative power density for each frequency band. In particular, the relative band power at each band was defined as the mean of the relative band power for each frequency bin within that band. Increasing tertiles values of alpha3/alpha2 frequency power ratio were estimated in all subjects and 2 groups were thus formed: low tertile (a3/a2<1.17;) and high tertile (a3/a2>1.17). The two groups of MCI has been demostrated in previous studies to be different in nature. In particular, the high alpha3/alpha 2 EEG power ratio MCI group is at major risk to convert to Alzheimer's disease, as well as to have different pattern of hippocampal atrophy, regional blood perfusion and basal ganglia and thalamus gray matter lesions as compared to the low alpha3/alpha2 power ratio MCI groups [23,32].

MRI scans

For each subject, a high-resolution sagittal T1 weighted volumetric MR scan was acquired at the Neuroradiology Unit of the 'Citta' di Brescia' Hospital, Brescia, by using a 1.0 T Philips Gyroscan scanner, with a gradient echo 3D technique: TR = 20 ms, TE = 5 ms, flip angle = 30, field of view = 220 mm, acquisition matrix 256 • 256, slice thickness 1.3 mm.

Cortical thickness estimation steps

Cortical thickness measurements for 74 MCI patients were made using a fully automated magnetic resonance imaging-based analysis technique: FreeSurfer, a set of software tools for the study of cortical and subcortical anatomy. Briefy, in the cortical surface stream, the models of the boundary between white matter and cortical gray matter as well as the pial surface were constructed. Once these surfaces are known, an array of anatomical measures becomes possible, including: cortical thickness, surface area, curvature, and surface normal at eash a gfeal meu5(si)4(5(aIisGf)8(e)4r)1-gnd snd stio(, a co)12(r)-10(t)-5(ic)-3(a)-5(l s)5(ur)-6(fa
as de sua ge(a)56(h)94(eT)19(o)78(dig)87(ha)19(5)(6)19(6)(t)6(3)(TBM)8(yS pan)3(s)-19(ang (ensGB)/MCI and 3surfaceEC)]TBM5 Span-21<75and-1.2(
orma84(b)7(le t)6emalate moden[35]e and(xtg)-5(rc(er)13(e)3bn)12((a)-5(lvg-8(oC xw)8(e)4(l)3(s a)9(r)13(e(r)13(mn)4(o)16vw)8(e)-5(e)6m.)]TEMC A
snb hasinelate(t)6(,de(t)6(et)-6minm)4(I t)-6(h)4(e(a)-3(a)9rg-8gw)8(-(s)-8ic)-3(a)-5(e, f)9(o75(d(in)8(I p)-5(a)19(t)10(t)6(et)-6en)8(y o)12(f t)-6(h)4(e
atrs(

alpha1 6.9–8.9 Hz; alpha2 8.9–10.9 Hz; alpha3 10.9–12.9 Hz. Finally, the relative power spectra was computed for each subject in the individual-determined frequency bands. The ratio between the absolute power and the mean power spectra from 2 to 45 Hz engenders the relative power density for each frequency band. In particular, the relative band power at each band was defined as the mean of the relative band power for each frequency bin within that band. Increasing tertiles values of alpha3/alpha2 frequency power ratio were estimated in all subjects and 2 groups were thus formed: low tertile (a3/a2<1.17;) and high tertile (a3/a2>1.17). The two groups of MCI has been demostrated in previous studies to be different in nature. In particular, the high alpha3/alpha 2 EEG power ratio MCI group is at major risk to convert to Alzheimer's disease, as well as to have different pattern of hippocampal atrophy, regional blood perfusion and basal ganglia and thalamus gray matter lesions as compared to the low alpha3/alpha2 power ratio MCI groups [23,32].

MRI scans

For each subject, a high-resolution sagittal T1 weighted volumetric MR scan was acquired at the Neuroradiology Unit of the 'Citta' di Brescia' Hospital, Brescia, by using a 1.0 T Philips Gyroscan scanner, with a gradient echo 3D technique: TR = 20 ms, TE = 5 ms, flip angle = 30, field of view = 220 mm, acquisition matrix 256 • 256, slice thickness 1.3 mm.

Cortical thickness estimation steps

Cortical thickness measurements for 74 MCI patients were made using a fully automated magnetic resonance imaging-based analysis technique: FreeSurfer, a set of software tools for the study of cortical and subcortical anatomy. Briefy, in the cortical surface stream, the models of the boundary between white matter and cortical gray matter as well as the pial surface were constructed. Once these surfaces are known, an array of anatomical measures becomes possible, including: cortical thickness, surface area, curvature, and surface normal at eash a gfeal meu5(si)4(5(aIisGf)8(e)4r)1-gnd snd stio(, a co)12(r)-10(t)-5(ic)-3(a)-5(l s)5(ur)-6(fa
as de sua ce(a)56(h)4(e T)19(o)78(d)-7(ha)19(5)6(1)9(6)10(t)6(1)TBM8(ySpan3(s)-1Lang (ensGB)/MCI and surfaceBC)]TBM5 TSpan-1<75 and1.2(
orma84(b)7(le t)6emalate moden[35]e and(xtg)-5(rc(er)13(e)3bn)12((a)-5(lvg-8(oC xw)8(e)4(l)3(s a)9(r)13(e(r)13(mn)4(o)16vw)8(e)-5(e)6m.)]TBMC
snb hasinelate(t)6(,de(t)6(et)-6minm)4(I t)-6(h)4(e(a)-3(a)9rg-8gw)8(-(s)-8ic)-3(a)-5(e, f)9(o75(d(in)8(I p)-5(a)19(t)10(t)6(et)-6en)8(y o)12(f t)-6(h)4(e
atrs(