

In Vitro Antimicrobial Activity of Crude Leaf Extracts from *Aloe secundiflora*, *Bulbine frutescens*, *Vernonia lasiopopus* and *Tagetes minuta* against *Salmonella typhi*

Rachuonyo HO^{1*}, Ogola PE², Arika WM², Nyamai DW² and Wambani JR³

¹Department of Microbiology, Kenyatta University, Kenya

²Department of Biochemistry and Biotechnology, Kenyatta University, Kenya

³Department of Medical Laboratory Sciences, Kenyatta University, Kenya

Abstract

Four medicinal plants leaves were investigated to evaluate their antibacterial potential of the methanol extracts against *Salmonella typhi* by disc diffusion method. The methanol extract from *Aloe secundiflora* showed strong antibacterial activity against a clinical isolate of *Salmonella typhi* at low concentrations (5.5 mg/ml) as compared to *Bulbine frutescens* extract (8.8 mg/ml). The minimum inhibitory concentration ranged from 5 mg/ml - 9 mg/ml whereas (15 µg/ml) was used as a positive control while dimethyl sulphoxide and distilled water were used as the negative control. The extracts were preliminary screened for the presence of secondary metabolites to determine the presence infections caused by *Salmonella typhi*.

*Corresponding author:

hibertrachuonyo@gmail.com

Received
15

Accepted February

Published February

Citation:

In Vitro

frutescens, *Vernonia lasiopopus* and *Tagetes minuta* against *Salmonella typhi*.

J

DOI: [10.4172/2573-4555.1000187](https://doi.org/10.4172/2573-4555.1000187)

whatman filter paper no. 1. The filtrate was poured into a round bottom flask and concentrated using a vacuum evaporator and stored in a labelled amber glass bottle at room temperature away from light and heat before being used for antibacterial efficacy test.

Antimicrobial susceptibility testing

The microorganism used was clinical isolate of *Staphylococcus aureus* obtained from Kenyatta University Health Centre Laboratory, Nairobi.

Conclusion

In conclusion, the present study will help us to use these medicinal plants as a source of herbal medicine to treat different infectious diseases caused by *S. aureus*, *E. coli*, and other gram negative bacterial pathogens. The extracts can also be used as raw material in the manufacturing of conventionally used drugs against pathogenic bacteria that have developed resistance against standard antibiotics. Further isolation of the specific bioactive compounds responsible for the antimicrobial activity will give us other natural resources that can be used in treating bacterial infections.

References

1. () The of a . British Journal .
Fabricant Farnsworth () The value of plants used in traditional medicine for drug discovery. Health 1 .
3. (1993) *Tagetes minuta* Herb from of the .