

## Increase in Colonoscopy Utilization Resulted in Risk of Colorectal Cancer

Kasem M\*

Department of Breast cancer, Syrian private university, UK

### Abstract

### Introduction

1. The prevalence of colorectal cancer (CRC) has increased significantly in many countries, including the United States, over the past few decades. This increase is largely attributed to the rising incidence of adenomatous polyps, which are precursors to CRC. The risk of developing CRC is also influenced by several factors, including diet, lifestyle, and genetics.

2. The introduction of colonoscopy as a screening tool for CRC has led to a significant increase in the detection of adenomatous polyps. This increase in detection has led to a corresponding increase in the risk of CRC, as more polyps are being identified and removed.

3. The risk of CRC is also influenced by the duration of colonoscopy. Studies have shown that the risk of CRC increases with the duration of colonoscopy, particularly in the first few years after the procedure.

4. The risk of CRC is also influenced by the quality of the colonoscopy. Studies have shown that the risk of CRC is higher in patients who undergo a low-quality colonoscopy compared to those who undergo a high-quality colonoscopy.

5. The risk of CRC is also influenced by the frequency of colonoscopy. Studies have shown that the risk of CRC is lower in patients who undergo colonoscopy more frequently compared to those who undergo it less frequently.

6. A. The risk of CRC is also influenced by the presence of adenomatous polyps. Studies have shown that the risk of CRC is higher in patients who have adenomatous polyps compared to those who do not have them.

7. A. The risk of CRC is also influenced by the presence of adenomatous polyps. Studies have shown that the risk of CRC is higher in patients who have adenomatous polyps compared to those who do not have them.

8. The risk of CRC is also influenced by the presence of adenomatous polyps. Studies have shown that the risk of CRC is higher in patients who have adenomatous polyps compared to those who do not have them.

### Acknowledgement

### Conflict of Interest

### References

sympathetic overactivity

\*Corresponding author: Kasem M, Department of Breast cancer, Syrian private

Received:

Revised:

Citation:

Copyright: ©

Editor assigned:

Reviewed:

Published:

0

Kasem M

