

+874A> IF - a ...
i i a a i a p n i a a n o m a o a i 18,19.B ,
i p o i p a p i p o i , a o i p
o a i p i n i a m o , m i n a
a n a i o n a i i a - o n i o i a
IF +874A/ o m o i o A A .
... p a a p o i p a a a .
a a a i p i p o a a o p a a 7, 2017.
o a i o a : (i p o R IF R i p
a m n a R i p o - R IF - R i p o - R IF -)

*Corresponding author: Shahab Alizadeh, Department of Cellular and Molecular Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences (TUMS), Tehran, Iran, Tel: +98-9197604090; Fax: +98-21-889 559 79; E-mail: sh_alizadeh@razi.tums.ac.ir

Received April 07, 2017; Accepted May 23, 2017; Published June 09, 2017

Citation: Razi B, Alizadeh S
evaluate the strength of association in the dominant model, recessive model and heterozygotes contrast.

Results: A total of 4 case-control studies, including 210 subjects, were included in this meta-analysis. Combined analysis of these studies showed that the polymorphism +874(T/A) and aplastic anemia risk in the overall population was OR=1.27, 95% CI=0.47-2.08, dominant model: OR=1.27, 95% CI=0.47-2.08, recessive model: OR=1.27, 95% CI=0.47-2.08, heterozygotes contrast: OR=1.27, 95% CI=0.47-2.08.

