

# Intravascular Lymphoma: Look for the Less Obvious!

Antonella Bianchi<sup>1\*</sup>, Lorenza Falcone<sup>1</sup>, Ombretta Annibaldi<sup>2</sup>, Lorenzo Ricci<sup>3</sup> and Mario Tombini<sup>3</sup>

\*Corresponding author: Antonella Bianchi, Unit of Pathology, University Hospital Campus Bio-Medico, Via Alvaro del Portillo, 200, Rome, Italy, Tel: +(39) 06.22541.1153; E-mail: a.bianchi@unicampus.it

Received date: March 29, 2018; Accepted date: April 24, 2018; Published date: May 02, 2018

Copyright: ©2018 Bianchi A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

Intravascular large B cell lymphoma (IVBCL) is a rare subtype of extranodal large B-cell lymphoma with a distinct presentation. Anatomically the disease is characterized by the proliferation of clonal lymphocytes within small vessels with relative sparing of the surrounding tissue. The clinical symptoms of the disease are dependent on the specific organ involvement, which often includes the central nervous system and skin. Because of the various modes of presentation and the rarity of IVBCL, the diagnosis is challenging and made post-mortem in most cases. The tumor is often disseminated at the time of diagnosis and prognosis is poor, even with aggressive chemotherapy. The spectrum of neurological presentations of IVBCL can be heterogeneous. We report a case of intravascular lymphoma diagnosed on autopsy.

**Keywords:** Lymphoma; Lymphoproliferative disease

**Abbreviations:** IVBCL: Intravascular Large B Cell Lymphoma

## Introduction

Intravascular large B cell lymphoma (IVBCL) is a rare type of non-Hodgkin's lymphoma (NHL) characterized by the proliferation of lymphoma cells within the lumina of small blood vessels, particularly capillaries and post-capillary venules, without an obvious extravascular tumor mass or readily observable circulating lymphoma cells in the peripheral blood. This type of lymphoma has also been described as intravascular lymphomatosis, angiotropic large cell lymphoma, and malignant angioendotheliomatosis. Originally described by Pfeiffer et al. in 1959 as an endothelial neoplasm with cutaneous involvement [1], this lymphoproliferative disease is now regarded as a well-defined entity in the WHO classification of lymphoid malignancy revised in 2017 [2]. The clinical presentation is varied and often includes symptoms related to organ dysfunction caused by occlusion of blood vessels. The diagnosis is challenging and, especially in the past, was usually recognized at the autopsy. In recent years, the heightened awareness of IVBCL has resulted in more patients being diagnosed during life, thereby allowing for treatment. The true incidence is unknown. Median age at diagnosis is in the sixth to seventh decades; there is no sex predilection [2].

There are two types of presentation: a Western form and an Asian variant, seen mainly in Japan. This lymphoma shows a propensity to manifest itself more frequently with CNS and cutaneous symptoms in Western countries or with hemophagocytic syndrome in Asian countries, mainly Japan [3,4].


Patients may be asymptomatic or symptomatic, presenting with fever of unknown origin, mental status, or personality changes.

## Case Presentation


An ex-smoker, diabetic, hypertensive 76-year-old man was admitted to our hospital with weight loss of about 10 kg in recent months, progressive cognitive decline, psycho-motor agitation, confusion and subsequent ideomotor slowdown in the context of a clinical history of relapsing ischemic stroke and worsening cognitive decay over a period of about 5 months. Some months before he had first episode of confusion, dysarthria, left facial deficiency of central type and postural instability, with a diagnosis of "cerebral ischemia in the frontal and right globus pallidus" and he began cardiovascular therapy with Clopidogrel. Later he underwent a new hospitalization for acute force deficit and drowsiness and subsequently discharged with the diagnosis of multiple ischemic lesions of the left and right hemisphere. During hospitalization in our hospital the brain CT scan revealed a low-density area in the cortical-subcortical right parietal lobe; laboratory tests showed elevated serum lactate dehydrogenase (LDH), beta-2 microglobulin, elevated sedimentation rate (VES) and elevated neuron specific enolase (NSE) and elevated antinuclear antibody (ANA). As cerebral vasculitis was suspected, corticosteroid therapy was initiated but clinical response was mild and transient.

Subsequently, clinical condition worsened completely and eventually

showed immunohistochemical stain for CD20 and were also documented in other tissue as lung kidney and heart (Figure 2).



**Figure 1:** Brain parenchyma softening (left) and evident vascular pattern (right).



**Figure 2** Atypical appearing lymphoid cells were observed within brain parenchyma small blood vessels (hematoxylin and eosin, original magnification 200X) (left) and were highlighted by positive immunoreactivity with CD20 (DAKO, original magnification 200X) (right).

## Discussion

The clinical diagnosis of IVBCL may be difficult [5]. Cases with cutaneous involvement can be confirmed by skin biopsy [6], but neurological symptoms can be heterogeneous and could be interpreted as vasculitis [7]. Delays in diagnosis due to the subtlety and focal nature of the intravascular infiltrates often lead to a terminal disease with death before chemotherapy is initiated [6].

Moreover, from the pathogenetic point of view, the precise mechanisms responsible for intravascular localization are at the moment largely unknown. Ponzoni et al. in an elderly but still valid immunohistochemical study proposed a very intriguing model. In a few words, in IVBCL, the second phase of the neoplastic cell-endothelial interaction, termed locking, is altered due to the lack on the surface of neoplastic cells of several molecules, as CD29 (beta integrin) and CD54 (ICAM-1), thus preventing movement of neoplastic cells through the vessels [1].

Other explanations have included the aberrant expression of CD11a and CD49d (VLA-4) on IVBCL cells as a possible mechanism because