Journal of
Alzheimer's Disease & Parkinsonism
ISSN: 2161-0460

Clement et al., Alzheimers Dis Parkinsonism 2016, 6:6
DOI: 10.4172/2161-0460.1000276

Research Article Open Access

noncoding and intronic regions of the loci. Exome sequencing may also miss exonic variants which are found close to exon/intron borders [8]. erefore these studies may have overlooked mutations which could a ect splicing through disrupting donor and acceptor splice sites.

Splicing plays an important role in human genetic disease. Almost a third of currently known disease-causing variants disrupt splicing, although this is likely to be an underestimate [9,10]. Aberrant splicing is implicated in many neurodegenerative disorders [11] and in LOAD, variants causing dysfunctional splicing have been found in some of the risk genes including PICALM and CD33 [12-14]. However the causative functional variant(s) within these loci are yet to be discovered. This study aimed to identify potential functional splicing mutations in the nine original GWAS-risk genes: CLU, PICALM, CR1, ABCA7, BIN1, the MS4A locus, CD2AP, EPHA1 and CD33. Target enriched next generation sequencing (NGS) was used to resequence the entire genetic region for each of these GWAS-risk loci in 96 LOAD patients and in silico splicing analysis and minigene assays were used to investigate the effect of novel variants identified. Two variants, rs881768 A>G in ABCA7 and a novel variant 11: 60179827 T>G in MS4A6A were shown, in these cell assays, to affect the splicing of these genes. The method employed in the paper successfully identified potential splicing variants in GWAS-risk genes. Further investigation will be needed to understand the full effect of these variants on LOAD risk. However, these results suggest a possible pipeline in order to identify putative functional variants as a result of NGS in disease-associated loci although improvements are needed within the current prediction programme in order to reduce the number of false positives.

Keywords: Late onset Alzheimer's disease; Next generation sequencing; Splicing; Annotation; Functional variations; Minigene assays

Introduction

Late onset Alzheimer's disease (LOAD) is an insidious neurodegenerative disease responsible for most cases of dementia in the elderly. Despite being widely studied, the disease still lacks a clear pathogenesis. However, there is a strong genetic component to LOAD with 21 disease-risk genetic regions having been identified through genome wide association studies (GWAS) [1]. Due to the nature of GWAS experimental design, the variants associated through GWAS are frequently not the causal functional variants generating the disease association. Instead, one on. Inste, CLU , PICALM, EPHA1, art from [5], have used targeted nts which may be found in the

*Corresponding author: Kevin Morgan, Human Genetics, School of Life Sciences, A Floor, West Block, Room 1306, Queens Medical Centre, University of Nottingham, Nottingham, NG7 2UH, United Kingdom, Tel: +44 115 8230724; E-mail: Kevin.Morgan@nottingham.ac.uk

Received October 03, 2016; Accepted October 22, 2016; Published October 29, 2016

Citation: Clement N, Braae A, Turton J, Lord J, Guetta-Baranes T, et al. (2016) Investigating Splicing Variants Uncovered by Next-Generation Sequencing the Alzheimer's Disease Candidate Genes, CLU, PICALM, CR1, ABCA7, BIN1, the MS4A Locus, CD2AP, EPHA1 and CD33. J Alzheimers Dis Parkinsonism 6: 276. doi: 10.4172/2161-0460.1000276

Copyright: © 2016 Clement N, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The genome analysis toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. Genome Res 20: 1297-1303.

23. McLaren W, Pritchard B, Rios D, Chen Y, Flicek P, et al. (2010) Deriving the consequences of genomic variants with the Ensembl API and SNP effect predictor. Bioinforma Oxf Engl 26: 2069-2070.

24. Consortium TEP (2011) A user's guide to the encyclopedia of DNA elements (ENCODE). PLoS Biol 9: e1001046.

25. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, et al. (2011) The variant