Institutionalized dependent older adults represent a highly vulnerable group of the population, with high mortality as described in previous studies [1,2]. e Activities of Daily Living (ADL) level [3], cognitive impairment [4], hip fracture [5], cancer [6] and Chronic Obstructive Pulmonary Disease [7] are important risk factors of mortality in this population. Recent studies have also found an increased risk of death among institutionalized older adults than those that live within the community, adjusting for age, gender and clinical conditions [8,9].

In Spain, elderly dependents receive care through the Public System of Social Services. During the six-year period 2007-2012, more than two hundred thousand older adults (200,000) had received care in Andalusia, the Spanish region with the greatest number of people registered in this system. rough this system, the population receive care in nursing homes or in their own homes, according to their own preferences and their social and clinical conditions [10].

e creation of a big data set allowed us to validate in our elderly population some predictors of mortality and con rm the importance of care setting as a risk of death factor. To do this, we use clinical and social data taken from the Information System for Dependence and the Electronic Health Record. e date of death is obtained from the Deaths Registry of Andalusia and the data is cross-referenced with that of the National Deaths Register. With them, we performed a survival analysis and included the following variables: sex, age, ADL level, preferences (nursing home or home-based care), support network characteristics (assessment, consistency, frailty and risk of claudication) and clinical conditions (dementia, Alzheimer's disease, cerebrovascular disease, Chronic Obstructive Pulmonary Disease, hip fracture and cancer).

We analyzed data from 200,039 people, 71% women. e average age was 80 years. According to the results, the risk of death is lower for women (38% less than men), and increases by ADL level (1,3% higher for each additional point on ADL Scale), age (5% for each additional year of age), Chronic Obstructive Pulmonary Disease (29%) and cancer (42%). A er adjusting these variables, we found that nursing home care increases 55% the risk of death compared to home-based care (Table 1).

Our results are consistent with previous studies. We found an elevated mortality of institutionalized older adults [2], and the greater risk of death in the residential care setting compared to home-based care [8,9]. e results also con rm that ADL level is a very powerful predictive mortality factor [3].

ese ndings highlight the importance of further understanding the mortality of the elderly in these two environments. Are there other variables that may help us explain this e ect? On the one hand, it is necessary to explore whether the geriatric assessment and health care programs act as protective factors for mortality of older adults that live in their own homes [11-13]. In Andalusia, there are speci c care plans for older adults in their homes such as the Telecare Service, the monitoring program a er hospital discharged, the follow-up visits to chronic patients by nursing personnel, health examinations for those aged 65 and above, etc. Another point to address is the di erential handling of chronic diseases in older people both within

***Corresponding author:** Sandra Arlette Pinzon, Professor, Andalusian School of Public Health, Citizenship and Social Welfare, Cuesta del Observatorio, 4 Campus Universitario de Cartuja Granada, Andalusia 18080, Spain, Tel: +34 677904299; E-mail: sandra.pinzon.easp@juntadeandalucia.es

Received September 28, 2016; **Accepted** October 03, 2016; **Published** October 10, 2016

Citation: Pinzon S, Rocha JL, Reyes V, Sanchez P, Martinez M, et al. (2016) Longevity: Big Data Comparison of Mortality in Older Adults Receiving Care in Nursing Homes or Their Own Home in Andalusia (Spain). J Alzheimers Dis Parkinsonism 6: 268. doi: 10.4172/2161-0460.1000268

Copyright: © 2016 Pinzon S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

16. Perkins MM, Ball MM, Whittington FJ, Hollingsworh C (2012) Relational autonomy in assisted living: A focus on diverse care settings for older adults. J Aging Stud 26: 214-225.

17. Skarsater I, Langius A, Agren H, Haggstrom L,