



© 2017

R

(R )

A A

A

A

A

A

A

A

C

C

References

1. Kusano-Tsunoh A, Nakatsuka H, Satoh H, Shimizu H, Sato S, et al. (2001) Effects of family-togetherness on the food selection by primary and junior high school students: family-togetherness means better food. *Tohoku J Exp Med* 194: 121-127.
2. Lubans D, Richards J, Hillman C, Faulkner G, Beauchamp M, et al. (2016) Physical activity for cognitive and mental health in youth: a systematic review of mechanisms. *Pediatrics* 138: 20161642.
3. Doi Y, Ishihara K, Uchiyama M (2014) Reliability of the strengths and difficulties questionnaire in Japanese preschool children aged 4-6 years. *J Epidemiol* 24: 514-518.
4. Matsuishi T, Nagano M, Araki Y, Tanaka Y, Iwasaki M, et al. (2008) Scale properties of the Japanese version of the Strengths and Difficulties Questionnaire (SDQ): a study of infant and school children in community samples. *Brain Dev* 30: 410-415.
5. Fulkerson JA, Story M, Mellin A, Lefert N, Neumark-Sztainer D, et al. (2006) Family dinner meal frequency and adolescent development: relationships with developmental assets and high-risk behaviors. *J Adolesc Health* 39: 337-345.
6. Eisenberg ME, Olson RE, Neumark-Sztainer D, Story M, Bearinger LH (2004) Correlations between family meals and psychosocial well-being among adolescents. *Arch Pediatr Adolesc Med* 158: 792-796.
- 7.