

Saloni Pahwa^{*}, Sharma Anila, Kamboj Meenakshi, Gupta Gurudutt and Pasricha Sunil

Department of Pathology, Rajiv Gandhi Cancer Institute and Research Centre, Delhi-110085, India

***Correspondence to:** Dr. Saloni Pahwa M.D., Department of Pathology, Rajiv Gandhi Cancer Institute and Research Centre, Delhi-110085, India, Tel: (+91)-799-611-9814; E-mail: pahwasaloni10@gmail.com

Received: December 07, 2020; **Accepted:** December 21, 2020; **Published:** December 28, 2020

Copyright: © 2020 Pahwa S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Gestational choriocarcinomas are malignant neoplasms generally arising in the uterus in women of child bearing age. These are aggressive tumours with high incidence of metastasis to vascular organs like lung, liver and brain. Renal metastasis is extremely rare with low incidence rate and very few cases reported in literature. Hereby, we report a rare case of metastatic choriocarcinoma to the kidney in a 29 year old female 10 years after resection of a Hydatidiform mole. The histopathological diagnosis was made on a nephrectomy specimen. Pelvic and abdominal

