[7]. Okra also stabilizes blood sugar levels and helps to control the rate at which sugar is absorbed in the body. In addition, okra mucilage has also been reported as an e ective tablet binder

w

e four varieties di ered signi cantly (<0.05) in 100 seed weight. While LD88 had the highest weight of 100 seeds (4.42 g), followed by NHAe-47-4 (4.25 g), V_{35} (4.22 g) the local variety had the least weight of a 100 seeds at 3.87 g. ese results are presented in Table 3.

D

e results of this study are indicative of the fact that the four varieties of okra evaluated were signi cantly di erent in some of the morphological traits but de nitely not in most of the yield attributes except for one hundred (100) seed weight. e observation of signi cant di erences in some of the traits is an indication that genetic diversity do exist among the varieties, thereby providing a basis for selection. is is in consonance with what has been reported earlier [21-23], where it was demonstrated that such genetic variability existed amongst okra varieties evaluated in the respective studies on genetic diversity among some okra germplasm.

In this study, the four varieties di ered signi cantly in the number of days to owering. It has been demonstrated that on a general basis, early owering is detrimental for overall productivity in okra as the source to sink ratio will be potentially limited for e ective photosynthesis [24]. Di erences in owering periods among the varieties in the current study imply that their maturity periods vary. Depending on the desire of the breeder or farmer, appropriate selection can thus be made for either early or late maturing plants.

Plant height of the varieties evaluated was also signi cantly di erent. e height of the plant can potentially a ect yield as those that are taller are usually more prone to windstorms in the event of heavy seasonal or monsoonal rains. Plant height at owering and fruiting are of particular interest for breeding programmes because the presence of plants with tall and thin stems will increase the rate of lodging near harvesting and this could lead to loss of dry matter and subsequent decrease in fruit yield. Number of days to bud emergence and plant height at maturity, among other agronomic characters, are some of the most variable traits that are necessary for selection programmes aimed at improving desirable traits in okra [25]. It is suggestive from this that number of days to and plant height at owering are controlled by the same genetic variables [26,27]. Consequently, selection for dwarf stature may thus be made on the local variety as it was shorter than the other three varieties evaluated in the current study (Table 3).

Variation was also observed in the number of leaves, pod length and one hundred seed weight of the cultivars. Since leaves serve as the sites for photosynthetic activities in any plant, an increase or a decrease in their number may have very serious implications for production of assimilates in the crop. Consequently, a greater number of them in any particular variety would be assumed to produce a better crop yield due to the higher photosynthetic capacity that is brought to bear by an increased leaf area index and a resultant higher fraction of intercepted radiation and its utilization e ciency [21]. e better performance of NHAe47-4 and V_{35} in the current study, amongst other reasons, can thus be attributed to the higher number of leaves as well as their larger leaf sizes (Table 3), which may have enabled these cultivars to produce greater assimilates during their photosynthetic activities. is position is reinforced by the suggestion that NHAe47-4 is a variety bred by NIHORT in Nigeria, which is characterized by early owering with thick fresh pods, short to medium plant height, deeply lobed leaves and profuse branching [28]. is suggestion is further supported when the published data is examined for the reasons behind the yield increases that have been reported for this variety. Strictly speaking, NHAe47-4 is very o en known to produce higher yield than most local and exotic varieties including V_{35} (an exotic variety that has the same morphological features as those of NHAe47-4), as it is well suited to the local environment in Nigeria. Also one hundred seed weight is usually associated with pod yield and other yield attributes in a seed bearing plant. An increased number of leaves and a higher seed weight would thus be important selection criteria in the improvement of okra accessions.

e tender green pod is o en considered as the most important and economical part of okra production since it is utilized as vegetable throughout the world. Consequently, fruit length in consonance with pod number and pod weight are the most important determinants of production or yield. us, selection based on these characters will be quite bene cial in okra breeding programmes, especially since three amongst the four cultivars evaluated in this study showed identical results for fruit length, which were statistically similar (Table 3). A similar pattern has been identi ed earlier, where it was reported that the Sabz pari variety of okra had the most promising result for maximum pod length in that study [29]. ese results are also in conformity with others reported earlier[25,30-33] , where statistically di erent results for pod length amongst the di erent okra cultivars evaluated were observed.

Overall, there were no signi cant variations in the number of days to seedling emergence, average number of pods per plant and average number of seeds per pod amongst the four okra varieties evaluated. is might be on account of the genetic characteristics of the cultivars. It is most probable that the genes controlling these character traits are dominant in all the varieties, which could have accounted for the absence of variation expressed amongst them. Consequently, these three traits cannot be considered as e ective strategies while formulating selection indices for the improvement of okra.

C

It is apparent on the basis of the ndings from this study that NHAe-47-4 and V_{35} okra varieties will be useful in breeding programmes where a higher crop yield is the dominant desire. Conversely, the local okra variety would be preferable where shorter and early maturing plants are desired since it had a comparatively shorter stature and an earlier owering date than NHAe-47-4, V_{35} or LD88. However, further studies need to be conducted with a wider range of local and exotic okra varieties to enable a more robust selection of useful accessions for breeding programmes, especially on yield improvement.

Refe ences

1 [illegible]