



Navigating Colposcopy: What Women Need to Know About the Procedure

Riant Adeli*

University of Zambia, Department of Obstetrics and Gynaecology, Zambia

Abstract

References

1. Herman JM, Chang DT, Goodman KA, Dholakia AS, Raman SP, et al. (2015) Phase 2 multi-institutional trial evaluating gemcitabine and stereotactic body radiotherapy for patients with locally advanced unresectable pancreatic adenocarcinoma. *Cancer* 121: 1128-1137.
2. Koong AC, Le QT, Ho A, Fong B, Fisher G, et al. (2004) Phase I study of stereotactic radiosurgery in patients with locally advanced pancreatic cancer. *Int J Radiat Oncol Biol Phys* 58: 1017-1021.
3. Koong AC, Christoferson E, Le QT, Goodman KA, Ho A, et al. (2005) Phase II study to assess the efficacy of conventionally fractionated radiotherapy followed by a stereotactic radiosurgery boost in patients with locally advanced pancreatic cancer. *Int J Radiat Oncol Biol Phys* 63: 320-323.
4. Didolkar MS, Coleman CW, Brenner MJ, Chu KU, Olexa N, et al. (2010) Image-guided stereotactic radiosurgery for locally advanced pancreatic adenocarcinoma results of first 85 patients. *J Gastrointest Surg* 14: 1547-1559.
5. Schellenberg D, Goodman KA, Lee F, Chang S, Kuo T, et al. (2008) Gemcitabine chemotherapy and single-fraction stereotactic body radiotherapy for locally advanced pancreatic cancer. *Int J Radiat Oncol Biol Phys* 72: 678-686.
- 6.