

NMOSD Management in the Setting of Chronic Hepatitis B and Latent Tuberculosis: A Case Report

Dylan Sadowsky BS1*, Kevin Delijani BS1, William Davis MD2, Amy Safadi MD3, Petra Brayo MD3 and Benjamin Osborne MD3


¹Georgetown University School of Medicine, Washington DC

²Department of Ophthalmology, Medstar Georgetown University Hospital, Washington DC

³Department of Neurology, Medstar Georgetown University Hospital, Washington DC

Abstract

Neuromyelitis Optica Spectrum Disorder (NMOSD) is an infammatory autoimmune disorder of the central nervous system, with optic neuritis and transverse myelitis as its most common presentations. Although immunomodulatory treatment options for NMOSD have expanded, preventing reactivation of latent infections in patients can still be a therapeutic challenge. Our patient presented to the emergency department during an NMOSD pseudo-relapse with untreated latent tuberculosis (TB) and chronic hepatitis B and was briefy treated with high-dose IV methylprednisolone. After confrmation that neither hepatitis B nor TB had reactivated, the patient was discharged on isoniazid and entecavir. A month later she was s



*Corresponding author: Dylan Sadowsky BS, Georgetown University School of Medicine, Washington DC, USA, Tel: 8633935622; E-mail: djs311@georgetown.edu

Received: 25-Jul-2022, Manuscript No. jceni-22-70155; Editor assigned: 27-Jul-2022, PreQC No. jceni-22-70155 (PQ); Reviewed: 01-Sep-2022, QC No. jceni-22-70155; Revised: 07-Sep-2022, Manuscript No. jceni-22-70155 (R); Published: 29-Sep-2022, DOI: 10.4172/jceni.1000158

Citation: Dylan Sadowsky BS, Kevin Delijani BS, William Davis MD, Amy Safadi MD, Petra Brayo MD, et al. (2022) NMOSD Management in the Setting of Chronic Hepatitis B and Latent Tuberculosis: A Case Report. J Clin Exp Neuroimmunol, 7: 158.

Page 2 of 2