Nutritional Management for Patients with Head and Neck Cancer: The Second Step of an Italian Survey: The Opinion of Italian Otolaryngologists

Marianna Trignani[1*], Melissa Laus[2], Valentina Mastronardi[2], Olga Leone[2], Marilina De Rosa[2], Giulio Campitelli[2], Angelo Di Pilla1, Giuseppe Santarelli[3], Albina Allajbej[1], Ambra Pamio[4], Domenico Genovesi[1] and Adelchi Croce[2]

[1]Department of Radiation Oncology, G. DAnnunzioUniversity of Chieti, SS. Annunziata Hospital, Chieti, Italy

[2]Department of Otorhinolaryngology, G. DAnnunzio University of Chieti, SS. Annunziata Hospital, Chieti, Italy

[3]Department of Nutrition, SS. Annunziata Hospital, Chieti, Italy

[4]Department of Hygiene and Public Health, G. DAnnunzio University, Chieti, Italy

[*]**Corresponding author:** Marianna Trignani, Department of Radiotherapy, SS Annunziata Hospital, Via Dei Vestini, 66100 Chieti, Italy, Tel: 3908713558244; Fax: 3908713557374; E-mail: marianna.trignani@unich.it

Rec date: January 13, 2016; **Acc date:** February 10, 2016; **Pub date:** February 16, 2016

Copyright: © 2016 Trignani M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: The purpose of this study was to survey the opinion of Italian otolaryngologist about nutritional management in patients with head and neck tumors.

Materials and Methods: A survey of 10 questions was e-mailed to 100 Italian centers of Otolaryngology.

Results: A total of 27 surveys were filled in. Nutritional supplementation in preventive phase was used by 37% respondents. The majority of respondents (88.3%) stated to use percutaneous endoscopic gastrostomy (PEG) in a reactive phase. Nutritional counselling before starting treatment was performed “rarely” by 33.3% of respondents while “always” by 29.6% of respondents; however 85.2% of respondents stated that medical nutritionist assessment should represent a standard procedure before starting an oncologic treatment.

Conclusions: Current practice about nutritional management for head and neck tumors is wide heterogeneous. Early or reactive approaches remain questionable, though otolaryngologists agree on the use of PEG in a reactive phase.

Keywords: Endoscopic gastrostomy; Survey; Nutrition; Head and neck cancer

Introduction

Growing evidence shows that more aggressive therapeutic regimens improve survival outcome of patients with head and neck cancers (HNC) [1]. ®neousɳ[e