

Nutritional Status and Menarcheal Age of Rural Adolescent Girls of Salboni Block of Paschim Medinipur, West Bengal, India

Kankana De*

Vidyasagar University, Medinipur, West Bengal, India

***Corresponding author:** Kankana De, Vidyasagar University, Medinipur, West Bengal, India, Tel: +919474714273; E-mail: dekankana@gmail.com

Received Date: October 5, 2016, **Accepted date:** October 15, 2016, **Published date:** October 25, 2016

Copyright: © 2016 De K. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Objectives of study:

- To study mean age at Menarche of studied population.
- To study whatever nutritional status effects mean age at menarche.

Design: This is a cross sectional study which is done on 1009 Adolescent girls who belong to Salboni block of district Paschim Medinipur of West Bengal, India. To study effect of nutrition on age at menarche, different anthropometric measurements are applied.

Setting: Data was collected from adolescent health clinic named Anwasha Clinic which is under Salboni Rural hospital, West Medinipur.

Participants: Participants are 10-19 years adolescent who are resident of Salboni (Rural) block, and belongs to agriculture based family, they are selected randomly for study.

Major outcomes: Data are bU € Ä € @ P € p ricu:

deficiency or imbalance of nutrients is of public health significance among adolescents all over the world.

signif cant with age (F= 17.3 P<0001), Percent body fat is statistically signif cant with Age (F= 38.7, P<0001).

Variable	10 years Mean(Sd)	11 years Mean(Sd)	12 years Mean(Sd)	13 years Mean(Sd)	14 years Mean(Sd)	15 years Mean(Sd)	16 years Mean(Sd)	17 years Mean(Sd)	18 years Mean(Sd)	19 years Mean(Sd)	F
BMI (kg/m ²)	18.28 (2.07)	19.02 (2.2)	18.58 (1.62)	18.96 (1.59)	19.65 (1.76)	19.86 (1.6)	20.05 (1.69)	20.02 (1.22)	20.16 (1.22)	20.17 (1.05)	17.3*
Percent Body Fat	18.17 (2.26)	20.21 (2.43)	20.4 (2.15)	20.63 (2.24)	21.18 (2.5)	21.41 (2.85)	22.23 (2.68)	22.41 (5.16)	22.66 (5.56)	24.04 (3.25)	38.76*

Significant difference P<0.001

Table 3 Anova table show age wise increase Anthropometric variables of Adolescents

Age	Height (cm)		Weight (kg)				BMI (Kg/sqm)		t
	Premenarcho	Post Menarcho	t	Premenarcho	Post Menarcho	t	Premenarcho	Postmenarcho	
10 years	147.45 (3.58)	148.34 (3.83)	1.08	37.46 (5.75)	43.42 (3.92)	6	16.93 (1.37)	19.53 (1.74)	8.3
11 years	145.88 (4.84)	146.13 (5.08)	0.25	36.01 (3.15)	41.71 (4.16)	7.7	16.89 (1.73)	19.66 (1.42)	21.3
12 years	151.37 (4.40)	153.32 (4.39)	1.54	37.38 (3.34)	45.75 (3.39)	6.8	16.62 (1.47)	19.25 (1.31)	7.81
13 years	148.61 (2.99)	150.68 (3.80)	1.81	37.00 (2.56)	44.16 (3.47)	6.9	16.76 (1.19)	19.44 (1.32)	6.62
14 years	148.65 (3.46)	150.12 (3.87)	1.65	36.72 (3.27)	43.39 (3.32)	8.1	15.87 (0.65)	19.98 (1.41)	8.31

Significant at P<0.0001

occurs between her age of 10 to 16 years, however; it shows a remarkable range of variation. General improvement in nutrition and health has been suggested to explain the downward trend. The earlier onset of menarche has also been seen in developing countries like Bangladesh [17].

Conclusion

The mean menarcheal age (determined by retrospective method) of the studied sample was 11.88 (1.23). This study had shown sexual maturation in early adolescence period in comparison with Eastern India girls and present study sample experiences their early menarche comparing to rest Indians. In global perspective the age at menarche in this study is less than most African and Asian studies but closer to most European and North American countries.

References

1. Onyiriuka AN, Egbagbe EE (2013) Anthropometry and Menarcheal Status of Adolescent Nigerian Urban Senior Secondary School Girls. *Int J Endocrinology Metab* 11: 71-5
2. Hochberg Z, Belski J (2013) Evo-devo of human adolescence: beyond disease models of early puberty. *BMC-Med* 11: 113
3. Sizonenko PC, Himes JH (1993) Introduction to the use and interpretation of anthropometry in adolescence. *Hormone Res* 39: 7-8
4. Heitmann BL (1990) Evaluation of body fat estimated from body mass index, skinfolds and impedance. A comparative study. *Eur J Clin Nutr* 44: 831-837.
5. Lifshitz F, Tarim O, Smith MM (1993) Nutrition in adolescence. *Endocr Metab Clinics North Am* 22: 673-83
6. Kusnita P, Damarati (2013) Pengaruh Status Gizi Remaja Terhadap Usia Menarche Pada Siswi SDN Dukuh Mananggal Surabaya. *J GDLHUB* 2: 28-31.
7. Mpora BO, Piloya T, Awor S, Ngwiri T, Laigong P, et al. (2014) Age at menarche in relation to nutritional status and critical life events among rural and urban secondary school girls in post-conflict Northern Uganda. *BMC Women's Health* 16: 44
8. Chumlea WC, Schubert CM, Roche AF, Kulin HE, Lee PA, et al. (2003) Age at menarche and racial comparison in US girls. *Pediatrics* 111: 110-113
9. Bose K, Mukhopadhyay A (2004) Nutritional status of adolescent Bengalee Boys. *Indian Paediatrics* 41: 633
10. Wyshak G, Frisch RE (1982) Evidence for a secular trend in age of menarche. *N Engl J Med* 306: 1033-1035
11. De K, Das S, Bose K, Chakraborty R (2013) Nutritional status of rural bengalee girls 10-18 years of Salboni, Paschim Medinipur, West Bengal, India. *Asian J Bio Life Sci* 2
- 12.