



# Obesity: Anesthetic and Caesarean Procedure Complications in the Teaching Centers of Cotonou

Annelie KH<sup>1\*</sup>, Eugène Z<sup>2</sup>, Hounkpatin B<sup>2</sup>, Véronique T<sup>3</sup>, Mewanou Serge<sup>4</sup>, Tchaou Blaise<sup>5</sup> and Chobli Martin<sup>6</sup>

1University Clinic of Endocrinology and Metabolic Diseases of Cotonou, University of Abomey-Calavi, Cotonou, Benin

2National Teaching Hospital of Mother and Child "Lagoon" (Cotonou), Benin

3University Clinical of Gynecology and Obstetrics of CNHU-HKM (Cotonou), Benin

4Department of Anesthesia, National Hospital Center Koutoukou Hubert Maga, University of Abomey-Calavi, Cotonou, Benin

5Department of Anesthesia and Reanimation, University of Parakou, Benin

6Department of Emergency, National Hospital Center Koutoukou Hubert Maga, University of Abomey-Calavi, Cotonou, Benin

**Presentation:** The dismalness townships of the heftiness at the ladies are the entanglements of pregnancy, the dystocia, and the labors by caesarian system. This study had for target to decide the recurrence of the heftiness at the parturient, to esteem the effect of the stoutness on the labor by cesarean and the confusions postoperative prompt in the showing medical clinics of Cotonou. **Patients and strategy:** We did an imminent, relative and diagnostic study of January 13 to April 13, 2012. 800 ninety-one continuous parturient having had a Cesarean was incorporate. We contemplated the sedative strategies, the procedures of control of the aeronautical ways, the quantity of fundamental tests for the tracheal intubation, the intricacies and aftercare of the initial 48 hours. A correlation has been made by the BMI. The weight has

pervasiveness of overweight of 35.3% and the one of the

has been arranged in two g

been recouped at 286 parturient is 32.10%. There were 703 (78.90%) Cesareans in crisis and 188 (21.09%) modified Cesareans. The stout parturient additionally required all the more often a few lumbar punctures (p As indicated by a WHO's report, in excess of 33% of the ladies and a fourth of the men in Africa are in overweight, and these extents should increment to 41% and 30% separately during the following ten years. The late inclinations in the urbanization of the creating nations and the internationalization of the food advertise add to change the conduct of individuals and their style of life. The changes of way of life, bound to the sustenance, progress of the conventional infant care practices, headed to the rise of the overweight and the corpulence [1,2]. A neighborly study prompted Cotonou restores a

