



Introduction

0.0 1 0 () 0.13 () 0.03 0.122% 0.13 -1.2 0.023



(13)

3 %

(3% / 11%)

30%

(11% / 10%)

2%

32

30%

3%

(1% - 10%)

3%

20

20

21

12%

10%

(/)

Risk of Death from Opioid Treatment

1, 00/ 2011, / 00/



1. Ralphs J, Williams A, Richardson P, Pither C, Nicholas M (1994) Opiate reduction in chronic pain patients: A comparison of patient-controlled reduction and staff controlled cocktail methods. *Pain* 79–88.
2. Rome J, Townsend C, Bruce B, Sletten C, Luedtke C (2004) Chronic noncancer pain rehabilitation with opioid withdrawal: Comparison of treatment outcomes based on opioid use status at admission. *Mayo Clin Proc* 759-68.
3. Thieme K, Gromnica-Ihle E, Flor H (2003) Operant behavioral treatment of fibromyalgia: A controlled study. *Arthritis Rheum.* 314-20.
4. Williams A, Richardson P, Nicholas M, Pither C, Harding V, Ridout K, et al. (1996) Inpatient vTw r.0004(Pain732Inpatientc gs/Fm0 D67to.9)-20(r B.yiodmiled)TJ

Acknowledgement

W

Conflict of Interest

W