

Outline of Low-Dose Gabapentin in a Pediatric Patient of Case Presentation

Dr. Clara Wang*

Abstract

Although rare, primary erythromelalgia can have a major negative impact on a person's quality of life. There is no accepted standard of care for erythromelalgia, despite the fact that many individuals with this ailment need systemic therapy. Here, we present the case of a 7-year-old girl who, following therapy with low-dose gabapentin had her erythromelalgia symptoms resolve without any negative side effects. We also go through the effectiveness and safety of using low-dose gabapentin to relieve pain in youngsters.





Table 1: Dosage of horizant for patients with restless legs syndrome in accordance with creatinine clearance.

Creatinine Clearance (mL/min)	Target Dose Regimen
>60	600 mg per day
30-59	Start at 300 mg per day and increase to 600 mg as needed
15-29	300 mg per day

*Corresponding author: Dr. Clara Wang, School of Medicine, University of California, Irvine, 1001 Health Sciences Rd, Irvine, CA 92697, 2022, QC No: jhcn-22-80925, Revised: 04-Nov-2022, Manuscript No: jhcn-22-80925(R), Published: 11-Nov-2022, DOI: 10.4172/jhcn.1000179

Citation: Wang C (2022) Outline of Low-Dose Gabapentin in a Pediatric Patient of Case Presentation. J Health Care Prev, 5: 179.

Copyright: © 2022 Wang C. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

