

## **Commentary**

Women with pregestational type 1 and type 2 diabetes mellitus continue to have poorer gestation issues than the background population, including a three-to fourfold advanced rate of perinatal mortality. Still, lower birth rates have lately been reported in centers involved in the UK National Gestation in Diabetes (NPID) 2015 inspection compared with those reported in the Confidential Enquiry

## References

1. pregnancy. *Diabetes care* 30(11): 2785-2791.
2. Lauenborg J, Mathiesen E, Ovesen P, Westergaard JG, Ekbom P, et al. (2003) Audit on stillbirths in women with pregestational type 1 diabetes. *Diabetes care* 26(5): 1385-1389.
3. Owens LA, Egan AM, Carmody L, Dunne F (2016) Ten years of optimizing outcomes for women with type 1 and type 2 diabetes in pregnancy—the Atlantic DIP experience. *J Clin Endocrinol Metab* 101(4): 1598-1605.
4. Murphy HR, Bell R, Cartwright C, Curnow P, Maresh M, et al. (2017) Improved pregnancy outcomes in women with type 1 and type 2 diabetes but substantial clinic-to-clinic variations: a prospective nationwide study. *Diabetologia* 60(9): 1668-1677.
5. de Valk HW, van Nieuwaal NH, Visser GH (2006) Pregnancy outcome in type 2 diabetes mellitus: a retrospective analysis from the Netherlands. *Rev Diabet Stud* 3(3): 134.

Changes